

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37



Clinical Practice Guidelines for Orthodontics and Dentofacial Orthopedics

This document may not be copied or reproduced without the
express written permission of the AAO

©American Association of Orthodontists
Created: 1996
Amended: 2001, 2009, 2010, 2012, 2014, 2016, 2017

TABLE OF CONTENTS

Introduction..... 3

Evidence-Based Dentistry..... 4

 Definition..... 4

 Levels of Evidence..... 4

 Evidence-Based Practice..... 4

Pretreatment Considerations..... 5

 Examination..... 5

 Diagnostic Records..... 5

 Referral..... 6

Diagnosis and Treatment Planning..... 6

Diagnostic and Treatment Considerations for Anomalies of Jaw Size, Relationship of Jaw to Cranial Base, Dental Arch Relationship and Dental Alveolus..... 7

Diagnostic and Treatment Considerations for Anomalies of Tooth Position, Discrepancies of Tooth Size, Arch Length and Arch Form..... 10

Diagnostic and Treatment Considerations for Abnormalities of the Dentition (number, size, and shape), Vitality, Eruption Pattern, and Periodontal Support..... 12

Diagnostic and Treatment Considerations for Dentofacial Functional Abnormalities..... 14

Orthodontic Considerations for Craniofacial Anomalies, Cleft Lip and Palate..... 16

Treatment Objectives and Limiting Factors..... 16

 Goals..... 17

 Limiting Factors..... 17

Treatment Consultation and Informed Consent..... 17

 Risks Associated with Orthodontic Treatment..... 18

Post Treatment Evaluation and Outcomes Assessment..... 19

 Post Treatment Records..... 19

 Positive Outcomes of Treatment..... 19

 Negative Outcomes of Treatment..... 19

Retention..... 20

Record Keeping..... 20

Transfer of Orthodontic Patients..... 20

 Recommendations to the Transferring Practitioner..... 20

 Recommendations to the Accepting Practitioner..... 21

Appendix A..... 22

 Historical Development..... 22

 Updating of Clinical Practice Guidelines..... 22

Appendix B: Selected References..... 23

Introduction

Orthodontics and Dentofacial Orthopedics is a specialty area of dentistry concerned with the supervision, guidance and correction of the growing or mature dentofacial structures, including those conditions that require movement of teeth or correction of malrelationships and malformations of their related structures and the adjustment of relationships between and among teeth and facial bones by the application of forces and/or the stimulation and redirection of functional forces within the craniofacial complex. Major responsibilities of orthodontic practice include the diagnosis, prevention, interception, and treatment of all forms of malocclusion of the teeth and associated alterations of their surrounding structures; the design, application, and control of functional and corrective appliances; and the guidance of the dentition and its supporting structures to attain and maintain optimal occlusal relations and physiologic and esthetic harmony among facial and cranial structures.

A specialist in orthodontics and dentofacial orthopedics meets educational standards established by the Commission on Dental Accreditation of the American Dental Association (ADA) and must possess advanced knowledge in biomedical, clinical, and basic sciences. This knowledge includes the biology of tooth movement, cephalometrics, orthodontic diagnosis, treatment planning, surgical orthodontics, biomechanical principles, the effects of growth and development on tooth movement, application of orthopedic forces to dentofacial structures, and patient management and motivation.

The American Association of Orthodontists (AAO) is the leading national organization of dentists who limit their practice to orthodontics and dentofacial orthopedics and is recognized by the ADA as the sponsoring organization of the national certifying board, the American Board of Orthodontics. The membership of the AAO includes the vast majority of practicing orthodontists in the United States and Canada. The AAO has the background, expertise, and professional responsibility to assist the dental profession and the public by developing clinical practice guidelines for orthodontics and dentofacial orthopedics. The AAO recognizes its role in upholding the public trust granted to it by presenting these clinical practice guidelines to help practitioners develop judgments on diagnosis, treatment planning, and timing of orthodontic and dentofacial orthopedic therapy. The primary concern of the AAO is the provision of high quality orthodontic care and the protection of the public.

Practice guidelines, as defined by the Institute of Medicine, are “systematically developed statements to assist practitioner and patient decisions about appropriate health care for specific clinical circumstances.”

The Orthodontic Clinical Practice Guidelines for Orthodontics and Dentofacial Orthopedics presented in this document are condition based and are related to the International Classification of Diseases, Clinical Modification, 10th Edition (ICD 10 codes). This approach recognizes the need for integrated treatment of oral and dentofacial conditions rather than isolated treatment procedures. These guidelines are also directed toward the process of patient care and outline considerations related to diagnosis, treatment, and quality of care.

These guidelines were derived from a professional consensus, based on a review of relevant clinical and scientific literature, the expert opinion of educators, and the clinical experience of practicing orthodontists. Similar documents written by other organizations and publications related to guideline development were also reviewed.

There are various professionally accepted philosophies regarding orthodontic diagnosis, treatment, and retention. Because of the nature of the doctor-patient relationship, the practitioner,

1 who is actively engaged in treating the patient, is in the best position to evaluate and interpret the
2 complexities, timing, and potential efficacy from among different philosophies and systems
3 available. Deviations from these guidelines may be appropriate based on professional judgment
4 and individual patient needs and preferences. Where a practitioner chooses to deviate from these
5 guidelines (based on the circumstances of a particular patient or for any other reason) the
6 practitioner is advised to note in the patient's record the reason for the procedure followed. Finally,
7 it should be understood that adherence to these guidelines does not guarantee a successful
8 treatment outcome.

9
10 The AAO recognizes that these guidelines may be used by insurance carriers and other payers,
11 attorneys in malpractice litigation, and various entities with an interest in orthodontics. The
12 Association encourages all interested persons to become familiar with the Guidelines. This
13 document was not developed to establish standards of care or to be used for reimbursement or
14 litigation purposes. The AAO cautions that these uses involve considerations that are beyond the
15 scope of the Guidelines.

16
17 The professional conduct of members of the AAO is governed by the Principles of Ethics and
18 Code of Professional Conduct of the AAO and the ADA.

19 20 **Evidence-Based Dentistry**

21 22 *Definition*

23
24 The following outline of orthodontic diagnostic and treatment considerations are evidence based
25 recommendations. Evidence-based dentistry (EBD) is an approach to oral health care that
26 requires the judicious integration of systematic assessments of clinically relevant scientific
27 evidence relating to the patient's oral and medical condition and history, with the dentist's clinical
28 expertise and the patient's treatment needs and preferences.

29 30 *Levels of Evidence*

31
32 Rating systems exist to evaluate the strength of various study designs. The Centre for Evidence-
33 based Medicine provides background information on this topic, as well as a commonly used table
34 for the "Levels of Evidence." In general, the levels of evidence, from strongest to weakest, are:

- 35
- 36 Meta-analysis
- 37 Systematic Review
- 38 Randomized Trial
- 39 Cohort Study
- 40 Case/Control Study
- 41 Case Series
- 42 Expert Opinion

43 44 *Evidence-Based Practice*

45
46 Evidence-based practice is assisted by critical evaluation of the body of literature on a specific
47 topic. In particular, well-conducted systematic reviews and meta-analyses can provide guidance
48 to assist orthodontists in clinical decision-making. Some resources for accessing evidence-based
49 literature are:

- 1 1. AAO Evidence Based Orthodontic Research Website: A collection of systematic
2 reviews, meta-analyses, practice guidelines, and summary statements on
3 orthodontic topics.
- 4
- 5 2. The ADA Center for Evidence-based Dentistry: A website which houses
6 information on evidence-based dentistry, as well as a listing of systematic reviews
7 in dentistry. Additionally, this site provides links to other evidence-based resources.
8
- 9 3. PubMed: PubMed comprises more than 25 million citations for biomedical literature
10 from MEDLINE, life science journals, and online books.
- 11
- 12 4. Cochrane Collaboration: An international nonprofit organization that develops
13 evidence-based systematic reviews on health care interventions.
14

15 **Pretreatment Considerations**

16
17 A screening examination may be performed to determine the nature of the orthodontic problem,
18 and to determine if and when treatment is indicated. When treatment is indicated, a
19 comprehensive examination must be performed that should include:
20

21 *Examination*

22 23 A. Chief Complaint

24 The chief complaint or the reason for seeking treatment should be recorded as described
25 by the patient, parent or legal guardian.
26

27 B. Medical and Dental History

28 An appropriate medical and dental history must be obtained as a part of the initial
29 evaluation of the patient. If treatment is to be delayed until a future date, an updated history
30 may be necessary. Patients/parents/legal guardians should be requested to advise the
31 orthodontist of any change in the patient's health history.
32

33 C. Clinical Examination

34 A comprehensive clinical examination should include the following with all findings
35 recorded in the patient's record:
36

- 37 1. An extraoral facial assessment to determine facial form, symmetry, soft-tissue
38 harmony, and status of the perioral musculature. This determines deviations from
39 normal regarding a patient's sagittal, vertical, and transverse maxillofacial
40 relationships and to assess the relationship of the dentition to the facial structures.
41
- 42 2. An intraoral examination to assess the condition of the hard and soft tissues of the
43 mouth, (including the periodontium) and the static and functional status of the
44 patient's occlusion.
45
- 46 3. An evaluation of the temporomandibular joint and associated musculature to
47 assess function and disease.
48
- 49 4. Verification of the presence of any oral parafunctional habits.
50

51 *Diagnostic Records*

1 Diagnostic records, along with a comprehensive examination and history, form the foundation
2 upon which a diagnosis and treatment plan with options are built, and are a standard of
3 orthodontic care.

4
5 Diagnostic records and tests will vary with the nature of the patient's condition but must be
6 sufficient to identify the problems, formulate a diagnosis, and allow the development of an
7 acceptable course of treatment goals. Where limited orthodontic procedures are anticipated,
8 diagnostic records may vary from those associated with comprehensive care. Limited or
9 comprehensive treatment encompasses all treatment techniques, including aligners or aligners in
10 combination with fixed appliances and auxiliaries to significantly alter the alignment or occlusion
11 and function. The gathering of appropriate diagnostic records should be considered a standard of
12 care to allow for proper diagnosis, treatment plan and treatment rendered.

13
14 Pretreatment unaltered diagnostic records for comprehensive orthodontic treatment should include
15 the following to establish a baseline for documenting pre-existing conditions, treatment and/or
16 growth changes:

- 17
18 1. Extraoral and intraoral images (may include digital or video images) to supplement
19 the clinical findings.
- 20
21 2. Dental casts (or digital models) to assess the inter-arch and intra-arch relationship
22 of the teeth, to help determine arch length and width requirements, and to assess
23 arch symmetry.
- 24
25 3. Intraoral and/or panoramic radiographs to assess the condition and developmental
26 status of the teeth and hard tissue supporting structures, and to identify any dental
27 anomalies or pathology.
- 28
29 4. Radiographic imaging to permit relative evaluation of the size, shape, and positions
30 of the relevant hard and soft tissue craniofacial structures including the dentition,
31 and to aid in the identification of skeletal anomalies and/or pathology. Three-
32 dimensional cone-beam computed tomography (CBCT) may be used as an imaging
33 source to obtain this information.

34 35 *Referral*

36
37 Practitioners must make a recommendation for referral of patients to general dentists, other dental
38 specialists, physicians, or other health care practitioners whenever, in the judgment of a
39 practitioner, referral would be in the best interest of a patient.

40 41 **Diagnosis and Treatment Planning**

42
43 Prior to the initiation of orthodontic treatment, a diagnosis of the patient's oral health condition
44 must be made. A diagnosis allows for the development of a treatment plan that addresses the
45 patient's chief complaint; medical and dental history, and dental, facial, skeletal, functional, and/or
46 psychosocial problems.

47
48 After a diagnosis has been established, a treatment plan must be developed. Such a plan will
49 facilitate coordination of the treatment objectives and the various methods available for addressing
50 them. A well-documented treatment plan should be based on the findings from the medical and
51 dental history, clinical examination, diagnostic records, a critical evaluation of the patient's needs

1 and preferences, and the clinician's professional judgement and preferences. A documented plan
2 should be a standard of care. The plan should include:

- 3
- 4 1. A list of the patient's dental, facial, skeletal, functional, and/or psychosocial
5 problems.
- 6
- 7 2. A diagnosis which coordinates the patient/parents/legal guardian's chief complaint
8 with the clinical findings.
- 9
- 10 3. A documented plan for therapy which includes treatment goals, appliance selection,
11 sequencing and timing of treatment, coordination with other health care providers,
12 and retention.
- 13

14 The treatment plan should be periodically reassessed throughout treatment with progress records
15 taken as deemed appropriate by the clinician. This reassessment should take into consideration
16 various limiting factors and establish short- and/or long-term objectives.

17 **Diagnostic and Treatment Considerations for Anomalies of Jaw Size, Relationship of Jaw** 18 **to Cranial Base, Dental Arch Relationship and Dental Alveolus**

19
20
21 The following conditions may indicate the need for orthodontic or dentofacial orthopedic treatment.
22 These conditions may be structural, functional or esthetic in nature and may appear in various
23 combinations, and are not limited to the outline below. Frequently used treatment options, which
24 may include the removal of primary or permanent teeth, are listed for each condition. Adjunctive
25 procedures to those listed used to supplement anchorage needs and improve treatment outcomes
26 include but are not limited to: osseointegrated implants, mini-screw implants, miniplates and other
27 temporary anchorage devices.

28 29 I. Maxillary/Dentoalveolar Hyperplasia (Large Maxilla)

30 31 A. Diagnostic Considerations

- 32
- 33 1. Anteroposterior
- 34 a. Mid-Face Protrusion
- 35 b. Dentoalveolar Protrusion
- 36 c. Distocclusion
- 37 d. Excess Overjet
- 38 e. Asymmetry
- 39
- 40 2. Vertical
- 41 a. Long, Lower Anterior Face Height
- 42 b. Maxillary Vertical Excess
- 43 c. Excessive Gingival Display
- 44 d. Deep Overbite
- 45 e. Open Bite
- 46 f. Lip Incompetency
- 47 g. Asymmetry
- 48
- 49 3. Transverse
- 50 a. Buccal Maxillary Cross-bite (unilateral or bilateral; functional or
51 structural)

- 1 b. Cant of the occlusal plane
- 2 c. Asymmetry
- 3
- 4 B. Treatment Options
- 5
- 6 1. Primary Dentition - Treatment Indicated Under Certain Circumstances,
- 7 Appliances Vary
- 8
- 9 2. Transitional Dentition
- 10 a. Functional/Orthopedic Appliances
- 11 b. Fixed or Removable Orthodontic Appliances
- 12 c. Space Maintenance
- 13
- 14 3. Adolescent Dentition
- 15 a. Functional/Orthopedic Appliances
- 16 b. Fixed or Removable Orthodontic Appliances
- 17 c. Fixed Orthodontic Appliances Adjunctive to Orthognathic Surgery
- 18 (surgery usually performed after majority of growth completed)
- 19
- 20 4. Adult Dentition
- 21 a. Fixed or Removable Orthodontic Appliances
- 22 b. Fixed Orthodontic Appliances Adjunctive to Orthognathic Surgery
- 23

24 II. Maxillary/Dentoalveolar Hypoplasia (Small Maxilla)

- 25
- 26 A. Diagnostic Considerations
- 27
- 28 1. Anteroposterior
- 29 a. Mid-Face Deficiency
- 30 b. Dentoalveolar Deficiency
- 31 c. Mesiocclusion
- 32 d. Anterior Cross-bite (functional or structural)
- 33 e. Negative Overjet
- 34 f. Asymmetry
- 35
- 36 2. Vertical
- 37 a. Short, Lower Anterior Face Height
- 38 b. Dentoalveolar Deficiency
- 39 c. Deep Overbite
- 40 d. Open Bite
- 41 e. Lip Redundancy
- 42 f. Asymmetry
- 43
- 44 3. Transverse
- 45 a. Lingual Posterior Cross-bite (unilateral or bilateral; functional or
- 46 structural)
- 47 b. Occlusal Plane Cant
- 48 c. Asymmetry
- 49 d. Transverse Deficiency without Posterior Cross-bite
- 50

51 B. Treatment Options

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

- 1. Primary Dentition
 - a. Functional/Orthopedic Appliance
 - b. Fixed or Removable Orthodontic Appliance
- 2. Transitional Dentition
 - a. Functional/Orthopedic Appliance
 - b. Fixed or Removable Orthodontic Appliance
- 3. Adolescent Dentition
 - a. Functional/Orthopedic Appliance
 - b. Fixed or Removable Orthodontic Appliance
- 4. Adult Dentition
 - a. Fixed or Removable Orthodontic Appliance
 - b. Fixed Orthodontic Appliance Adjunctive to Orthognathic Surgery

III. Mandibular/Dentoalveolar Hyperplasia (Large Mandible)

A. Diagnostic Considerations

- 1. Anteroposterior
 - a. Prognathic Facial Pattern
 - b. Mesioocclusion
 - c. Anterior Cross-bite (functional or structural)
 - d. Macrogenia
 - e. Asymmetry
- 2. Vertical
 - a. Open Bite
 - b. Deep Overbite
 - c. Long Lower Facial Height/Steep Mandibular Plane Angle
 - d. Asymmetry
- 3. Transverse
 - a. Posterior Cross-bite (unilateral or bilateral; functional or structural)
 - b. Asymmetry

B. Treatment Options

- 1. Primary Dentition - Treatment Indicated Under Certain Circumstances, Appliances Vary
- 2. Transitional Dentition
 - a. Functional/Orthopedic Appliance
 - b. Fixed or Removable Orthodontic Appliance
- 3. Adolescent Dentition
 - a. Functional/Orthopedic Appliance
 - b. Fixed or Removable Orthodontic Appliance
- 4. Adult Dentition

- 1 a. Fixed or Removable Orthodontic Appliance
- 2 b. Fixed Orthodontic Appliance Adjunctive to Orthognathic Surgery
- 3
- 4 IV. Mandibular/Dentoalveolar Hypoplasia (Small Mandible)
- 5
- 6 A. Diagnostic Considerations
- 7
- 8 1. Anteroposterior
- 9 a. Mandibular Retrognathic Facial Pattern
- 10 b. Excess Overjet
- 11 c. Distoclusion
- 12 d. Asymmetry
- 13
- 14 2. Vertical
- 15 a. Open Bite
- 16 b. Deep Overbite
- 17 c. Short Lower Face Height
- 18 d. Long Lower Face Height
- 19
- 20 3. Transverse
- 21 a. Posterior Cross-bite (unilateral or bilateral; functional or structural)
- 22 b. Asymmetry
- 23
- 24 B. Treatment Options
- 25
- 26 1. Primary Dentition - Functional/Orthopedic Appliance
- 27
- 28 2. Transitional Dentition
- 29 a. Functional/Orthopedic Appliance
- 30 b. Fixed or Removable Orthodontic Appliance
- 31
- 32 3. Adolescent Dentition
- 33 a. Functional/Orthopedic Appliance
- 34 b. Fixed or Removable Orthodontic Appliance
- 35 c. Fixed Orthodontic Appliance Adjunctive to Orthognathic Surgery
- 36 (surgery usually performed after majority of growth completed)
- 37
- 38 4. Adult Dentition
- 39 a. Fixed or Removable Orthodontic Appliance
- 40 b. Fixed Orthodontic Appliance Adjunctive to Orthognathic Surgery
- 41

42 **Diagnostic and Treatment Considerations for Anomalies of Tooth Position, Discrepancies**
 43 **of Tooth Size, Arch Length, and Arch Form**

44
 45 These conditions may appear in various combinations and are not limited to the following.
 46 Frequently used treatment options, which may include the removal of primary or permanent teeth,
 47 are listed for each condition. Adjunctive procedures to those listed include modification of tooth
 48 size, restorative replacement, surgical exposure, and appropriate soft tissue surgery.

- 49
- 50 I. Deficient Arch Length (Crowding)
- 51

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

- A. Diagnostic Considerations
 - 1. Facial-Lingual Displacement
 - 2. Supra/Infra Eruption
 - 3. Rotations
 - 4. Impactions
 - 5. Axial Inclination of Teeth (Anterior or Posterior)
 - 6. Tooth Size
 - 7. Premature Loss of Primary Teeth
 - 8. Ankylosis
 - 9. Supernumeraries and aplasias
 - 10. Frenal attachments
 - 11. Transpositions

- B. Treatment Options
 - 1. Primary Dentition
 - a. Fixed or Removable Space Maintainer
 - b. Extraction of primary teeth

 - 2. Transitional Dentition
 - a. Functional/Orthopedic Appliance
 - b. Fixed or Removable Orthodontic Appliance
 - c. Serial Extraction

 - 3. Adolescent Dentition
 - a. Fixed or Removable Orthodontic Appliance
 - b. Functional/Orthopedic Appliance
 - c. Extractions of Permanent or Remaining Primary Teeth

 - 4. Adult Dentition
 - a. Fixed or Removable Orthodontic Appliance
 - b. Extraction of Permanent Teeth

 - 5. Interdisciplinary Referral

II. Excessive Arch Length (Spacing)

- A. Diagnostic Considerations
 - 1. Skeletal Arch Size
 - 2. Tooth Size
 - 3. Congenitally Missing Teeth
 - 4. Supernumeraries and Aplasias
 - 5. Axial Inclination of Teeth
 - 6. Facial-Lingual Displacement
 - 7. Rotations
 - 8. Fibrous Gingival Hyperplasia
 - 9. Frenal Attachments

- B. Treatment Options

1. Primary Dentition - Treatment Rarely Indicated
2. Transitional Dentition - Fixed or Removable Orthodontic Appliance
3. Adolescent Dentition - Fixed or Removable Orthodontic Appliance
4. Adult Dentition - Fixed or Removable Orthodontic Appliance
5. Interdisciplinary Referral

III. Discrepancies of Arch Form

A. Diagnostic Considerations

1. Asymmetry
2. Interarch Coordination
3. Abnormal Occlusal Planes: Curves of Wilson and Spee
4. Bi-level Occlusal Plane

B. Treatment Options

1. Primary Dentition - Fixed or Removable Orthodontic Appliance
2. Mixed Dentition
 - a. Fixed or Removable Orthodontic Appliance
 - b. Functional/Orthopedic Appliance
3. Adolescent Dentition
 - a. Fixed or Removable Orthodontic Appliance
 - b. Functional/Orthopedic Appliance
4. Adult Dentition
 - a. Fixed or Removable Orthodontic Appliance
 - b. Fixed Orthodontic Appliance Adjunctive to Orthognathic Surgery

Diagnostic and Treatment Considerations for Abnormalities of the Dentition (number, size, and shape), Vitality, Eruption Pattern, and Periodontal Support

Anomalies of tooth number, morphology or eruption pattern should be diagnosed and managed as soon as reasonably practical according to the particular requirements of each clinical situation. These conditions may appear in various combinations, and may indicate the need for orthodontic or dentofacial orthopedic treatment. Some of the frequently used treatment options may require a multidisciplinary approach and may include the extraction of primary or permanent teeth.

A. Diagnostic Considerations

1. Supernumerary Teeth
2. Missing Teeth
 - a. Congenital (Anodontia)
 - b. Pathologic
 - c. Traumatic
 - d. Extracted
3. Ectopic Erupting Teeth
4. Impacted Teeth
5. Eruption Anomalies

- 1 6. Over-Retained Primary Teeth
- 2 7. Ankylosed Teeth
- 3 8. Transposition
- 4 9. Atypical Crown Morphology
- 5 10. Premature Loss of Primary Teeth
- 6 11. Atypical Root Morphology
- 7 12. Root Resorption
- 8 13. Carious or Fractured Teeth
- 9 14. Character of Hard and Soft Tissue Supporting Structures
- 10 15. Tooth Vitality

11
12 **B. Treatment Options**

- 13
- 14 1. Supernumerary Teeth
- 15 a. Surgical Intervention
- 16 b. Extraction
- 17 c. Fixed or Removable Orthodontic Appliance
- 18 d. No Treatment
- 19
- 20 2. Missing Teeth
- 21 a. Space Maintenance/Space Regaining
- 22 b. Prosthetic Replacement of Teeth/Implants
- 23 c. Transplantation
- 24 d. Maintenance of Primary Teeth
- 25 e. Space Closure
- 26 f. Fixed or Removable Orthodontic Appliance
- 27
- 28 3. Ectopic Teeth
- 29 a. Extraction
- 30 b. Surgical Intervention
- 31 c. Fixed or Removable Orthodontic Appliance
- 32
- 33 4. Impacted Teeth
- 34 a. Surgical Intervention
- 35 b. Extraction
- 36 c. Fixed or Removable Orthodontic Appliance
- 37 d. No Treatment
- 38
- 39 5. Eruption Anomalies
- 40 a. Surgical Intervention
- 41 b. Retention with or without Coronal Modification
- 42 c. Extraction
- 43 d. Fixed or Removable Orthodontic Appliance
- 44 e. Referral for Medical Evaluation
- 45
- 46 6. Over-Retained Primary Teeth
- 47 a. Extraction
- 48
- 49 7. Ankylosed Teeth
- 50 a. Extraction
- 51 b. Surgical Luxation and/or Repositioning

- 1 c. Fixed or Removable Orthodontic Appliance
- 2 d. Retention with or without Coronal Modification
- 3
- 4 8. Transposition
- 5 a. Extraction
- 6 b. Retention with or without Coronal Modification
- 7 c. Transplantation
- 8 d. Fixed or Removable Orthodontic Appliance
- 9
- 10 9. Atypical Tooth Morphology
- 11 a. Retention with or without Coronal Modification
- 12 b. Extraction
- 13 c. Fixed or Removable Orthodontic Appliance
- 14
- 15 10. Premature Loss of Primary Teeth
- 16 a. Space Maintenance
- 17 b. Fixed or Removable Orthodontic Appliance
- 18
- 19 11. Atypical Root Morphology
- 20 a. Monitor Radiographically
- 21 b. Extraction
- 22
- 23 12. Root Resorption
- 24 a. Monitor Radiographically
- 25 b. Extraction
- 26 c. Stabilization
- 27 d. Treatment Alternative of Initiating Rest Periods
- 28
- 29 13. Carious or Fractured Teeth
- 30 a. Reposition Tooth or Root
- 31 b. Monitor Radiographically
- 32 c. Extraction
- 33 d. Fixed or Removable Orthodontic Appliance
- 34

Diagnostic and Treatment Considerations for Dentofacial Functional Abnormalities

Dentofacial functional abnormalities may occur in combination with other dentofacial conditions and should be diagnosed and managed according to the particular requirements of each clinical situation. Correction or control of functional problems may involve alteration of behavior patterns, may require orthodontic/dentofacial orthopedic treatment, or multidisciplinary treatment. The influence of functional abnormalities on dentofacial development is variable, and cause and effect relationships are difficult to determine.

- 44 A. Diagnostic Considerations
- 45
- 46 1. Lip Size and Function
- 47
- 48 2. Tongue Size and Function
- 49 a. Abnormal Tongue Function
- 50 b. Ankyloglossia
- 51 c. Microglossia or Macroglossia

- 1 3. Deleterious Habits
- 2 a. Thumb, Finger or Lip Sucking
- 3 b. Pacifier Sucking
- 4 c. Tongue Thrust/Sucking
- 5 d. Clenching
- 6 e. Clenching and Bruxism
- 7 f. Lip/Cheek Biting
- 8 g. Nail Biting
- 9 h. Foreign Objects (e.g., pipes, pens, pencils, musical instruments)
- 10 i. Smoking and/or Drug Usage
- 11
- 12 4. Airway Obstruction
- 13 a. Nasopharyngeal Morphology
- 14 b. Sleep Apnea
- 15 c. Allergies
- 16 d. Pathology
- 17
- 18 5. Speech Disorders
- 19
- 20 6. Mandibular Dysfunction
- 21 a. Dental Interferences
- 22 b. Skeletal Abnormalities
- 23 c. Neuromuscular Abnormalities
- 24 d. Temporomandibular Dysfunction
- 25
- 26 7. Trauma
- 27
- 28 8. Temporomandibular Disorders
- 29 Temporomandibular disorders represent a broad range of conditions which involve
- 30 medical, dental, and psychological factors. Such disorders may be associated with
- 31 stress, habits, emotional disorders, structural malrelationships, oro-facial pain,
- 32 trauma to the face or head, occlusal disharmonies, and medical problems
- 33 associated with osteoarthritis, rheumatoid arthritis, or viral disease. These factors
- 34 may be associated with temporomandibular disorders in one individual with no
- 35 symptomatology or pathology in another.
- 36

37 B. Treatment Options

- 38
- 39 1. Lip Size and Function
- 40 a. Fixed or Removable Orthodontic Appliance
- 41 b. Therapeutic Exercises/Myofunctional Therapy
- 42 c. Functional/Orthopedic Appliance
- 43 d. Surgery
- 44
- 45 2. Tongue Size and Function
- 46 a. Fixed or Removable Orthodontic Appliance
- 47 b. Therapeutic Exercises/Myofunctional Therapy
- 48 c. Functional/Orthopedic Appliance
- 49 d. Surgical Reduction
- 50 e. Lingual Frenectomy
- 51

- 1 3. Deleterious Habits
- 2 a. Behavior Management
- 3 b. Functional/Orthopedic Appliance
- 4 c. Therapeutic Exercises
- 5 d. Fixed or Removable Orthodontic Appliance
- 6
- 7 4. Airway Obstruction
- 8 a. Referral for Evaluation/Treatment/Surgery
- 9 b. Functional/Orthopedic Appliance
- 10 c. Orthognathic Surgery
- 11
- 12 5. Speech Disorders
- 13 a. Fixed or Removable Orthodontic Appliance
- 14 b. Referral for Evaluation/Treatment/Myofunctional Therapy
- 15
- 16 6. Mandibular Dysfunction
- 17 a. Occlusal Equilibration (Modification of Tooth Form)
- 18 b. Fixed or Removable Orthodontic Appliance
- 19 c. Fixed Orthodontic Appliance Adjunctive to Surgery
- 20 d. Functional/Orthopedic Appliance
- 21
- 22 7. Temporomandibular Disorders
- 23 Numerous treatment modalities, including orthodontics, have produced beneficial
- 24 results in the management of temporomandibular disorders. However, no singular
- 25 treatment modality may necessarily be definitive for any particular patient. There is
- 26 no scientific proof that any particular method of orthodontic treatment, whether
- 27 involving extraction or non-extraction, has any causative effect on
- 28 temporomandibular disorders. There is no reliable method for predicting or
- 29 preventing future temporomandibular disorders in any particular individual.
- 30

31 **Orthodontic Considerations for Craniofacial Anomalies, Cleft Lip and Palate**

32
33 Management of patients with these and other anomalies is, in many cases, best provided by a
34 multidisciplinary team of dentists and physicians and other healthcare professionals. The optimal
35 time for the first evaluation of these patients is within the first few days of life, and referral for team
36 evaluation and management is appropriate at any age. Treatment plans should be developed and
37 implemented on the basis of team recommendations. The orthodontist, as a member of the
38 craniofacial team, should obtain baseline diagnostic records, assist in treatment planning, and
39 perform orthodontic treatment as needed taking into account those factors that may influence
40 surgical management of the patient.

41
42 For patients at risk for developing malocclusion or maxillomandibular discrepancy, diagnostic
43 records should be collected at appropriate intervals. Depending on the goals to be accomplished,
44 alternating periods of treatment and retention may be necessary beginning at birth. For example,
45 patients with cleft lip and cleft palate may require presurgical maxillary orthopedics to improve the
46 position of the maxillary alveolar segments prior to lip and palate closure. Later in life, timing of
47 bone grafting of alveolar clefts to unify the segments should be determined by the stage of dental
48 development and with the collaboration of the orthodontist and surgeon.

49 **Treatment Objectives and Limiting Factors**

50
51

1 **Goals**

2
3 The goals of orthodontic treatment are optimum dentofacial function, health, stability and esthetics.
4 While these goals are desirable, it should be recognized that individual patients have problems,
5 concerns and conditions which may prevent the attainment of optimal results in every case, and
6 that the non-attainment of some of the goals of orthodontic treatment in a particular patient is no
7 indication of negligence by the orthodontist even when no limiting factors are present.
8

9 **Limiting Factors**

10
11 Orthodontic treatment results may be affected by extenuating circumstances beyond the
12 practitioner's control. These limiting factors should be documented in the patient's record when
13 they occur and the patient/parent/guardian should be informed. The following are some of the
14 more common limiting factors affecting orthodontic therapy:
15

- 16 1. Severity of the pretreatment condition
- 17 2. Pretreatment agreement to pursue limited objectives
- 18 3. Abnormal skeletal morphology or growth, during or after treatment
- 19 4. Abnormal size, shape, or number of teeth
- 20 5. Aberrant tooth eruption patterns
- 21 6. Patient's failure to initiate timely treatment, continue or complete treatment
- 22 7. Compromised periodontal tissues
- 23 8. Persistent deleterious habits or abnormalities of muscle function relating to the
- 24 dentofacial complex
- 25 9. Inability or unwillingness of the patient to cooperate with treatment (e.g., the wear
- 26 and/or care of appliances, oral hygiene measures, diet, or keeping appointments)
- 27 10. Failure to complete all recommended aspects of treatment
- 28 11. Poor quality, untimely or inappropriate integration of other recommended or
- 29 required dental and/or medical services
- 30 12. Medical complications or underlying systemic conditions
- 31 13. Patient transferring to or from another provider during orthodontic treatment
- 32 14. Incomplete correction or relapse of orthognathic surgical procedures
33

34 **Treatment Consultation and Informed Consent**

35
36 A discussion must be held with the patient/parents/legal guardian utilizing lay terminology to
37 provide sufficient information for the responsible party to accept or reject the proposed treatment
38 plan. This discussion must be documented and should include:
39

- 40 1. A description of the diagnosis and treatment plan.
- 41 2. A discussion of reasonable alternative treatments.
- 42 3. The relevant risks, compromises, and limitations associated with the proposed
- 43 treatment plan and alternative treatments.
- 44 4. A discussion of any portion of the treatment plan that will require the services of
- 45 other dental or medical health care providers and the anticipated effects of such
- 46 services on the orthodontic treatment plan.
- 47 5. The prognosis related to all treatment plans, including the option of no treatment.
- 48 6. A discussion of the patient's responsibility relating to the care (e.g., maintaining
- 49 periodic recall visits with their general dentist).
- 50 7. An estimate of the duration of active treatment and retention.

- 1 8. A signed agreement regarding informed consent and the financial arrangements
2 may be considered.

3 4 *Risks Associated with Orthodontic Treatment*

5
6 All forms of medical and dental treatment, including orthodontics, involve some risks and/or
7 limitations. Fortunately, in orthodontics, serious complications are infrequent. The orthodontist
8 should determine which potential risks to disclose to the patient in the exercise of sound
9 professional judgment given the clinical condition of the patient. Due to the length of orthodontic
10 treatment, conditions may arise which are coincident, but not caused by orthodontic treatment.
11 Some of the risks associated with orthodontic treatment include:

- 12
13 1. Tooth decay, or permanent markings (decalcification).
- 14 2. The length of the roots of teeth may become shortened. This may be of no clinical
15 significance or may require the discontinuance of orthodontic treatment with
16 subsequent interdisciplinary treatment to stabilize the teeth. In some cases root
17 shortening may be pre-existing.
- 18 3. The health of the bone and periodontal support of the teeth may be affected.
- 19 4. The teeth and/or jaws have a tendency to change their positions after treatment.
- 20 5. Temporomandibular joint problems may appear concurrently with orthodontic
21 treatment, but may not be related to the treatment.
- 22 6. The vitality of a tooth may be compromised.
- 23 7. Orthodontic appliances may irritate or damage the oral tissues and may cause
24 injury if accidentally swallowed or aspirated.
- 25 8. Dental materials, instruments, and equipment may result in damage or injury to the
26 oral tissues, face and/or eyes.
- 27 9. Accidents during treatment or patient misuse of orthodontic appliances may result
28 in injury to the oral tissues, face and/or eyes.
- 29 10. Oral surgery, orthognathic surgery or other adjunctive medical, surgical or dental
30 procedures may be recommended and/or necessary in conjunction with orthodontic
31 treatment. Associated treatments carry additional risks, limitations and additional
32 informed consent issues which must be discussed with the patient/parents/legal
33 guardian by the health care practitioner providing the service.
- 34 11. Orthodontic appliances may cause attrition, flaking or fracturing of tooth structure.
- 35 12. When orthodontic appliances are removed, fracture and/or damage to the teeth
36 may result.
- 37 13. Medical or psychosocial conditions may result in compromised results or
38 dissatisfaction with treatment.
- 39 14. Orthodontic materials may cause allergic reactions in some individuals.
- 40 15. Patients may be dissatisfied with their dental or facial esthetics at the conclusion of
41 treatment due to unrealistic expectations or perceptions.
- 42 16. Abnormal growth during or after treatment may produce undesirable results.
- 43 17. Treatment time may be extended and results compromised due to unforeseen
44 circumstances and poor patient cooperation.
- 45 18. Tooth movement during orthodontics may be adversely affected for patients
46 receiving certain pharmaceuticals as they have the potential to slow tooth
47 movement and may lengthen treatment time. The effects of these medications may
48 be severe enough to stop tooth movement which may result in removal of
49 appliances regardless of tooth positions. The effects of certain pharmaceuticals on
50 an individual are not predictable.

- 1 19. The use of orally applied drugs, especially certain drugs of abuse such as cocaine
2 or amphetamines, may seriously compromise the gums and bone tissue around
3 teeth which can be exacerbated by orthodontic treatment.
4

5 **Post Treatment Evaluation and Outcomes Assessment**

6
7 The effects of orthodontic treatment should be evaluated retrospectively with reference to the
8 pretreatment condition. Consistent re-evaluation of treatment results along with continued review
9 of treatment modalities and their effectiveness will serve to provide the public with the highest
10 quality of orthodontic care. Assessments of the outcome of treatment are dependent in part upon
11 the treatment goals and objectives, the condition being treated, the stage of the patient's
12 dentofacial development, the treatment provided and the patient's compliance as well as tissue
13 response to the therapy performed. Limiting factors must be considered when evaluating
14 treatment and outcomes.
15

16 *Post Treatment Records*

17
18 Post treatment unaltered records provide information for the quantitative and qualitative
19 assessment of treatment changes as well as for education, research, and quality assurance. Post
20 treatment records may include, but are not limited to:

- 21
22 1. Extra and intraoral images (digital, still or video images)
23 2. Dental casts (hard copy or digital format)
24 3. Intraoral, panoramic, and/or radiographic imaging to permit relative evaluation of
25 the size, shape, and positions of the relevant hard and soft tissue craniofacial
26 structures including the dentition.
27 4. Other indicated procedures or tests
28

29 *Positive Outcomes of Treatment*

- 30
31 1. Satisfaction of the patient's chief complaint
32 2. Well aligned teeth
33 3. Good or improved occlusal function
34 4. Good or improved dental and facial esthetics
35 5. Good or improved environment for dentofacial development
36 6. Desirable modification of the size, shape, and position of the jaw(s)
37 7. Stability of the treatment results
38 8. Good or improved dental and periodontal health
39

40 *Negative Outcomes of Treatment*

- 41
42 1. The patient's chief complaint was not satisfied
43 2. Poorly aligned teeth
44 3. Poor or unimproved occlusal function
45 4. Poor or unimproved dental and facial esthetics
46 5. Premature root resorption (primary teeth)
47 6. Excessive root resorption (permanent teeth)
48 7. Loss of periodontal support
49 8. Clinically significant decalcification or dental caries
50 9. Unsatisfactory modification of the size, shape, and position of the jaws
51 10. Instability of the treatment results

Retention

1. A retention plan must be established after reviewing the patient's original condition, treatment objectives, the results achieved, and/or any limiting factors.
2. Completion of orthodontic treatment does not ensure the stability of the result. Future treatment may be recommended when post treatment changes occur, which may be due to growth, maturation, aging, lack of compliance with the retention protocol, periodontal problems, oral habits and post treatment trauma, among other factors.
3. Post treatment changes may be minimized with an indefinite retention wear protocol.

Record Keeping

The keeping and preserving of a patient's dental record is necessary to the goal of providing high quality orthodontic treatment. Prudent record keeping is the foundation for planning and maintaining the continuity of patient care. It also provides documentary evidence of the evaluation and diagnosis of the patient's condition, the treatment plan, the treatment provided, referrals made, and follow up care. It also documents communications with the patient, other health care providers and any other third parties. The dental record also protects the legal interests of all parties. In addition, a patient's dental record may provide material for continuing education, research, administrative oversight, billing, and quality assurance.

1. Treatment procedures, changes in the treatment plan, patient compliance, treatment difficulties, and other important aspects of treatment must be recorded and maintained. Copies of related correspondence and appropriate release forms must also be maintained as part of the patient's record.
2. Documentation must be written, dictated, or computer annotated; maintained concurrently; and kept chronologically with any changes conspicuously noted.
3. The original records are usually considered the property of the practitioner. Laws regarding patient record access, duplication and transfer vary from state to state. Practitioners can obtain clarification from their state regulatory agency.
4. Electronic/digital records have the potential to be altered. Alteration of original electronic/digital records must be avoided. Credible computer software either prevents this or records any alteration of an original electronic/digital record. However, enhancement of images is allowed as long as these are duly labeled and saved as separate images. Enhancement of other electronic/digital records, such as radiographs, to enable better identification of landmarks and/or dentoskeletal anomalies is permissible; however, the original cannot be altered. It is the responsibility of the practitioner to protect the sanctity of all patient records as prescribed by all local, state and federal laws.

Transfer of Orthodontic Patients

Because of the time required to complete orthodontic treatment, the transfer of care from one practitioner to another is a common occurrence.

Recommendations to the Transferring Practitioner

1. Practitioners should attempt to arrange for the continuation of orthodontic treatment of their patients with as little interruption as possible. Regardless of the reason for

1 transfer, reasonable efforts of both the transferring and accepting practitioner are
2 necessary to effect an orderly transfer. It is recommended, and in some states
3 required, to obtain a written release from the patient/parents/legal guardian prior to
4 the transfer of the patient's records. It is preferable to send copies of the pertinent
5 records directly to the new practitioner. The use of electronic media may facilitate
6 this process. It is acceptable, but less desirable, to provide these records to the
7 patient/ parents/legal guardian. A patient's records should not be withheld due to an
8 outstanding balance.

- 9 2. The transferring practitioner should ensure that all appliances are in good order.
10 The patient/parents/legal guardian should be advised that extended periods of
11 active orthodontic treatment without supervision can be detrimental, and an
12 appointment with the new practitioner should be scheduled as soon as possible.
- 13 3. The patient/parents/legal guardian should be informed that there may be different
14 approaches to treatment by different practitioners.
- 15 4. The patient/parents/legal guardian should be informed that there may be different
16 fees with treatment by different practitioners.
- 17 5. The transferring practitioner should make no statements that would undermine the
18 establishment of a sound doctor-patient relationship with the accepting practitioner.
- 19 6. The transferring practitioner should be available for consultation by the accepting
20 practitioner.
- 21 7. The transferring practitioner should provide appropriate financial information in
22 advance or immediately upon request to the accepting practitioner.

23 24 *Recommendations to the Accepting Practitioner*

- 25
26 1. The accepting practitioner should review the patient's records, including the
27 previous financial arrangements if available, prior to the development of a plan for
28 continuation of treatment. In addition, the estimated time required to complete
29 treatment and the financial arrangement for continuation of treatment should be
30 discussed as soon as possible. Patients should be informed about their present oral
31 health status without unprofessional comments about prior treatment.
- 32 2. Appropriate records documenting the status of the case at the time of transfer
33 should be made.
- 34 3. A practitioner is not obligated to accept an orthodontic transfer patient. If a
35 practitioner is unable or unwilling to accept the transfer patient, the practitioner may
36 assist the patient/parents/legal guardian in finding another practitioner.
- 37 4. At the patient/parents/legal guardian's request, a practitioner may remove
38 appliances from a patient not of record. If appropriate, previous practitioners
39 should be consulted.

40
41 Members should be aware of the following documents written by the AAO Legal Counsel:

- 42
43 1. [Second Opinions](#)
- 44 2. [Terminating the Doctor/Patient Relationship](#)
- 45 3. [Patient Records and Record Keeping](#)

1 **Appendix A**

2

3 **Historical Development**

4

5 At its November 1993 meeting, the AAO Board of Trustees directed the AAO Council on
6 Orthodontic Health Care (COHC) to study the feasibility of developing clinical practice guidelines
7 for orthodontics. The council met in January 1994 and proposed a business plan for the
8 development of Guidelines, which was considered at the February 1994 meeting of the AAO
9 Board of Trustees. It was the consensus of the AAO Board of Trustees to develop guidelines
10 utilizing the expertise within the AAO. A task force was appointed.

11

12 The task force met three times between July 1994 and January 1995 and wrote draft guidelines. A
13 copy of draft guidelines was sent to all active AAO members in April 1995 for review. Open forums
14 were held at the 1995 AAO Annual Session and at the meetings of all eight AAO constituent
15 societies during August-November 1995. The task force met again in December 1995 to revise the
16 draft guidelines based on feedback received in 1995. The December 1995 revised draft guidelines
17 were widely circulated in January 1996 for comment. The task force reviewed the comments and a
18 revised draft of the guidelines was distributed to the AAO House of Delegates members, the Board
19 of Trustees and other leaders of organized orthodontics in April 1996. An open forum was held at
20 the 1996 AAO Annual Session for comments on the revised draft guidelines. The revised draft
21 guidelines were approved by the Board of Trustees, a House of Delegates Reference Committee
22 and by the House of Delegates. The Clinical Practice Guidelines were printed in 1996 and were
23 made available to AAO members.

24

25 **Updating of Clinical Practice Guidelines**

26

27 The American Association of Orthodontists considers its Clinical Practice Guidelines to be a living
28 document. The existence of this document is intended to stimulate improvement in the practice of
29 orthodontics by identifying areas where knowledge is incomplete or inadequate. The AAO
30 recognizes the dynamic nature of orthodontics and dentofacial orthopedics and the necessity for
31 updating the guidelines to reflect the evolving science and art of orthodontics. Revisions to the
32 document, with opportunities for AAO member input, will occur periodically.

1 **Appendix B**

2
3 **Selected References**

4
5 **Introduction**

6
7 American Association of Orthodontists. American Association of Orthodontists bylaws and
8 principles of ethics. St. Louis: American Association of Orthodontists; 1994, amended through
9 2015.

10
11 American Association of Orthodontists. Glossary of dentofacial orthopedic terms. St. Louis:
12 American Association of Orthodontists; 2012.

13
14 American Dental Association. Council on Ethics, Bylaws and Judicial Affairs. Principles of ethics
15 and code of professional conduct with official advisory opinions revised to April 2012., Chicago:
16 American Dental Association; 2012.

17
18 American Dental Association. Standards for advanced specialty education programs in
19 orthodontics. Chicago: American Dental Association; 1991.

20
21 American Dental Association. Commission on Dental Accreditation. Accreditation Standards for
22 Advanced Specialty Education Programs in Orthodontics and Dentofacial Orthopedics. Chicago:
23 American Dental Association; 2013.

24
25 Daskalogiannakis, J. Glossary of orthodontic terms. Chicago: Quintessence Publishing Co., Inc.;
26 2000.

27
28 **Evidence-Based Dentistry**

29
30 American Association of Orthodontists. Council on Scientific Affairs. Evidence Based Orthodontic
31 Research Resource Center. Available at:
32 <https://www.aaoinfo.org/library-research/resource-centers/evidence-based-orthodontic-research>.
33 Accessed October 27, 2015.

34
35 American Dental Association. Center for Evidence-Based Dentistry. Available at:
36 <http://ebd.ada.org/en/about>. Accessed October 27, 2015.

37
38 American Dental Association. Center for Evidence-Based Dentistry. Available at:
39 <http://ebd.ada.org>. Accessed October 27, 2015.

40
41 The Cochrane Collaboration. Cochrane Database of Systematic Reviews. Available at:
42 <http://www.cochrane.org>. Accessed October 27, 2015.

43
44 Howick J, Chalmers I, Glasziou P, Greenhalgh T, Heneghan C, Liberati A, Moschetti I, Phillips B,
45 Thornton H. The 2011 Oxford CEBM evidence levels of evidence (introductory document). Oxford
46 Centre for Evidence-Based Medicine. Available at:
47 <http://www.cebm.net/wp-content/uploads/2014/06/CEBM-Levels-of-Evidence-Introduction-2.1.pdf>.
48 Accessed October 27, 2015.

1 Howick J, Chalmers I, Glasziou P, Greenhalgh T, Heneghan C, Liberati A, Moschetti I, Phillips B,
2 Thornton H, Goddard O, Hodgkinson M. The Oxford 2011 Levels of Evidence. Oxford Centre for
3 Evidence-Based Medicine. Available at:
4 <http://www.cebm.net/wp-content/uploads/2014/06/CEBM-Levels-of-Evidence-2.1.pdf>. Accessed
5 October 27, 2015.

6
7 Huang GJ, Richmond S, Vig KWL. Evidence-based orthodontics, Chichester, West Sussex, UK:
8 Wiley-Blackwell; 2011.

9
10 US National Library of Medicine. National Institutes of Health. PUBMED. Available at:
11 <http://www.ncbi.nlm.nih.gov/pubmed>. Accessed October 27, 2015.

12 13 **Pretreatment Considerations**

14
15 Ackerman JL, Nguyen T, Proffit WR. Diagnosis and treatment planning in orthodontics. In: Graber
16 LW, Vanarsdall RL, Vig KWL. Orthodontics: current principles and techniques. 5th ed.
17 Philadelphia: Elsevier/Mosby; 2012. p. 3-58.

18
19 Ackerman JL, Proffit WR. The characteristics of malocclusion: a modern approach to classification
20 and diagnosis. American Journal of Orthodontics 1969;56:443-54.

21
22 Albino JE. Psychosocial aspects of malocclusion. In: Matarazzo JT, Miller NE, Weiss SM, Herd
23 JA, Weiss SM. Behavior health: a handbook of health enhancement and disease prevention. New
24 York: Wiley; 1984. p. 918-29.

25
26 American Board of Orthodontics. Case Record Preparation. Available at:
27 [https://americanboardortho.com/orthodontic-professionals/about-board-certification/clinical-](https://americanboardortho.com/orthodontic-professionals/about-board-certification/clinical-examination/case-record-preparation)
28 [examination/case-record-preparation](https://americanboardortho.com/orthodontic-professionals/about-board-certification/clinical-examination/case-record-preparation). Accessed January 18, 2016.

29
30 American Dental Association. Council on Scientific Affairs. U.S. Department of Health and Human
31 Services. Food and Drug Administration. Dental radiographic examinations: recommendations for
32 patient selection and limiting radiation exposure. Chicago: American Dental Association; 2012.

33
34 Assaf AT, Zrnc TA, Remus CC, Schönfeld M, Habermann CR, Riecke B, Friedrich RE, Fiehler J,
35 Heiland M, Sedlacik J. Evaluation of four different optimized magnetic-resonance-imaging
36 sequences for visualization of dental and maxillo-mandibular structures at 3 T. Journal of Cranio
37 Maxillofacial Surgery 2014;42:1356-63.

38
39 Ballrick JW, Fields HW, Beck M, Sun Z, Germak J. The cervical vertebrae staging method's
40 reliability in detecting pre and post mandibular growth. Orthodontic Waves 2013;72:105-11.

41
42 Bjerklin K, Ericson S. How a computerized tomography examination changed the treatment plans
43 of 80 children with retained and ectopically positioned maxillary canines. Angle Orthodontist
44 2006;76:43-51.

45
46 Burstone CJ. Application of bioengineering to clinical orthodontics. In: Graber LW, Vanarsdall RL,
47 Vig KWL. Orthodontics: current principles and techniques. 5th ed. Philadelphia: Elsevier/Mosby;
48 2012. p. 345-380.

1 Cartsos VM, Zhu S, Zavras AI. Bisphosphonate use and the risk of adverse jaw outcomes: a
2 medical claims study of 714,217 people. *Journal of the American Dental Association* 2008;139:23-
3 30.
4
5 Dale JG, Dale HC. Interceptive guidance of occlusion, with emphasis on diagnosis. In: Graber LW,
6 Vanarsdall RL, Vig KWL. *Orthodontics: current principles and techniques*. 5th ed. Philadelphia:
7 Elsevier/Mosby; 2012. p. 423-476.
8
9 Devereux L, Moles D, Cunningham SJ, McKnight M. How important are lateral cephalometric
10 radiographs in orthodontic treatment planning? *American Journal of Orthodontics and Dentofacial*
11 *Orthopedics* 2011;139:e175-81.
12
13 Eichenberger M, Baumgartner S. The impact of rapid palatal expansion on children's general
14 health: a literature review. *European Journal of Paediatric Dentistry* 2014;15:67-71.
15
16 Eley KA, Watt-Smith SR, Golding SJ. "Black bone" MRI: a potential alternative to CT when
17 imaging the head and neck: report of eight clinical cases and review of the Oxford experience.
18 *British Journal of Radiology* 2012;85:1457-64.
19
20 Eley KA, Watt-Smith SR, Golding SJ. "Black Bone" MRI: a potential non-ionizing method for three-
21 dimensional cephalometric analysis – a preliminary feasibility study. *Dento maxillo facial Radiology*
22 2013;42:20130236.
23
24 Ellis PE, Benson PE. Does articulating study casts make a difference to treatment planning?
25 *Journal of Orthodontics* 2003;30:45-9.
26
27 European Commission, Nuclear Energy Directorate, Radiation Protection. Cone beam CT for
28 dental and maxillofacial radiology : evidence-based guidelines (Radiation protection ; no 172).
29 Luxembourg : Directorate-General for Energy, 2012. Available at
30 http://www.sedentext.eu/files/radiation_protection_172.pdf. Accessed November 5, 2015.
31
32 Fleming PS, Marinho V, Johal A. Orthodontic measurements on digital study models compared
33 with plaster models: a systematic review. *Orthodontics and Craniofacial Research* 2011;14:1-16.
34
35 Flores-Mir C, Korayem M, Heo G, Witmans M, Major MP, Major PW. Craniofacial morphological
36 characteristics in children with obstructive sleep apnea syndrome: a systematic review and meta-
37 analysis. *Journal of the American Dental Association* 2013;144:269-77.
38
39 Gabriel DB, Southard KA, Qian F, Marshall SD, Franciscus RG, Southard TE. Cervical vertebrae
40 maturation method: poor reproducibility. *American Journal of Orthodontics and Dentofacial*
41 *Orthopedics* 2009;136:478. e1-7; discussion 478-80.
42
43 Halazonetis DJ. Cone-beam computed tomography is not the imaging technique of choice for
44 comprehensive orthodontic assessment. *American Journal of Orthodontics and Dentofacial*
45 *Orthopedics* 2012;141:403, 405, 407.
46
47 Haney E, Gansky SA, Lee JS, Johnson E, Maki K, Miller AJ, Huang JC. Comparative analysis of
48 traditional radiographs and cone-beam computed tomography volumetric images in the diagnosis
49 and treatment planning of maxillary impacted canines. *American Journal of Orthodontics and*
50 *Dentofacial Orthopedics* 2010;137:590-7.
51

1 Hartsfield JK. Personalized orthodontics: limitations and possibilities in orthodontic practice. In:
2 Krishnan V, Davidovitch Z. Biological mechanisms of tooth movement. 2nd ed. Chichester, West
3 Sussex ; John Wiley & Sons Inc.; 2015. p. 164-172.
4
5 Hou HM, Hägg U, Sam K, Rabie AB, Wong RW, Lam B, Ip, MS. Dentofacial characteristics of
6 Chinese obstructive sleep apnea patients in relation to obesity and severity. *Angle Orthodontist*
7 2006;76:962-9.
8
9 Isaac A, Major M, Witmans M, Alrajhi Y, Flores-Mir C, Major P, Alsufyani N, Korayem M, El-Hakim
10 H. Correlations between acoustic rhinometry, subjective symptoms, and endoscopic findings in
11 symptomatic children with nasal obstruction. *JAMA Otolaryngology - Head and Neck Surgery*
12 2015;141:550-5.
13
14 Krieger E, Jacobs C, Walter C, Wehrbein H. Current state of orthodontic patients under
15 bisphosphonate therapy. *Head and Face Medicine* 2013;9:10.
16
17 Larson BE. Cone-beam computed tomography is the imaging technique of choice for
18 comprehensive orthodontic assessment. *American Journal of Orthodontics and Dentofacial*
19 *Orthopedics* 2012;141:402, 404, 406.
20
21 Leifert MF, Leifert MM, Efstratiadis SS, Cangialosi TJ. Comparison of space analysis evaluations
22 with digital models and plaster dental casts. *American Journal of Orthodontics and Dentofacial*
23 *Orthopedics* 2009;136:16.e1-4, discussion 16.
24
25 Mortensen MG, Kiyak HA, Omnell L. Patient and parent understanding of informed consent in
26 orthodontics. *American Journal of Orthodontics and Dentofacial Orthopedics* 2003;124:541-50.
27
28 Moyers RE, van der Linden FPGM, Riolo ML, McNamara JA. Standards of human occlusal
29 development. Craniofacial growth series No. 5 Ann Arbor, MI: Center for Human Growth and
30 Development, University of Michigan; 1976.
31
32 Neeley WW, Kluemper GT, Hays LR. Psychiatry in orthodontics. Part 1: Typical adolescent
33 psychiatric disorders and their relevance to orthodontic practice. *American Journal of Orthodontics*
34 *and Dentofacial Orthopedics* 2006;129:176-84.
35
36 Nestman TS, Marshall SD, Qian F, Holton N, Franciscus RG, Southard TE. Cervical vertebrae
37 maturation method morphologic criteria: poor reproducibility. *American Journal of Orthodontics*
38 *and Dentofacial Orthopedics* 2011;140:182-8.
39
40 Nijkamp PG, Habets LL, Aartman IH, Zentner A. The influence of cephalometrics on orthodontic
41 treatment planning. *European Journal of Orthodontics* 2008;30:630-5.
42
43 O'Reilly RL, O'Riordan JW, Greenwood AM. Orthodontic abnormalities in patients with eating
44 disorders. *International Dental Journal* 1991;41:212-6.
45
46 Pae EK, McKenna GA, Sheehan TJ, Garcia R, Kuhlberg A, Nanda R. Role of lateral
47 cephalograms in assessing severity and difficulty of orthodontic cases. *American Journal of*
48 *Orthodontics and Dentofacial Orthopedics* 2001;120:254-62.
49
50 Rheude B, Sadowsky PL, Ferriera A, Jacobson A. An evaluation of the use of digital study models
51 in orthodontic diagnosis and treatment planning. *Angle Orthodontist* 2005;75:300-4.

1 Riolo ML, Moyers RE, McNamara JA Jr, Hunter WS. An Atlas of craniofacial growth. Craniofacial
2 growth series No. 2 Ann Arbor, MI: Center for Human Growth and Development, University of
3 Michigan; 1974.
4
5 Rischen RJ, Breuning KH, Bronkhorst EM, Kuijpers-Jagtman AM. Records needed for orthodontic
6 diagnosis and treatment planning: a systematic review. PLoS ONE 2013;8:e74186.
7
8 Sadeghianrizi A, Forsberg CM, Marcus C, Dahllöf G. Craniofacial development in obese
9 adolescents. European Journal of Orthodontics 2005;27:550-5.
10
11 Silva MA, Wolf U, Heinicke F, Bumann A, Visser H, Hirsch E. Cone-beam computed tomography
12 for routine orthodontic treatment planning: a radiation dose evaluation. American Journal of
13 Orthodontics and Dentofacial Orthopedics 2008;133:640.e1-5.
14
15 Simmons KE. Electronic medical record and its implications for orthodontists. American Journal of
16 Orthodontics and Dentofacial Orthopedics 2011;139:567-8.
17
18 Stevens DR, Flores-Mir C, Nebbe B, Raboud DW, Heo G, Major PW. Validity, reliability, and
19 reproducibility of plaster vs digital study models: comparison of peer assessment rating and Bolton
20 analysis and their constituent measurements. American Journal of Orthodontics and Dentofacial
21 Orthopedics 2006;129:794-803.
22
23 van der Linden FPGM, Duterloo, HS. Development of the human dentition an atlas. Hagerstown,
24 MD: Harper & Row; 1976. Also available as ebook with 50 video clips exclusively from Apple
25 iBooks.
26
27 van Vlijmen OJ, Kuijpers MA, Bergé SJ, Schols JG, Maal TJ, Breuning H, Kuijpers-Jagtman AM.
28 Evidence supporting the use of cone-beam computed tomography in orthodontics. Journal of the
29 American Dental Association 2012;143:241-52.
30
31 Vanarsdall RL, Musich DR. Adult interdisciplinary therapy: diagnosis and treatment. In: Graber
32 LW, Vanarsdall RL, Vig KWL. Orthodontics: current principles and techniques. 5th ed.
33 Philadelphia: Elsevier/Mosby; 2012. p. 843-896.
34
35 Whetten JL, Williamson PC, Heo G, Varnhagen C, Major PW. Variations in orthodontic treatment
36 planning decisions of Class II patients between virtual 3-dimensional models and traditional plaster
37 study models. American Journal of Orthodontics and Dentofacial Orthopedics 2006;130:485-91.
38
39 Will L. Psychological aspects of orthodontics. In: Graber LW, Vanarsdall RL, Vig KWL.
40 Orthodontics: current principles and techniques. 5th ed. Philadelphia: Elsevier/Mosby; 2012. p. 99-
41 108.
42
43 **Diagnosis and Treatment Planning**
44
45 Ackerman JL, Nguyen T, Proffit WR. The decision making process in orthodontics. In: Graber LW,
46 Vanarsdall RL, Vig KWL. Orthodontics: current principles and techniques. 5th ed. Philadelphia:
47 Elsevier/Mosby; 2012. p. 3-58.
48
49 Andreasen JO, Paulsen HU, Yu Z, Ahlquist R, Bayer T, Schwartz O. A long-term study of 370
50 autotransplanted premolars. Part I. Surgical procedures and standardized techniques for
51 monitoring healing. European Journal of Orthodontics 1990;12:3-13.

1 Andreasen JO, Paulsen HU, Yu Z, Bayer T, Schwartz O. A long-term study of 370
2 autotransplanted premolars. Part II. Tooth survival and pulp healing subsequent to transplantation.
3 European Journal of Orthodontics 1990;12:14-24.
4
5 Andreasen JO, Paulsen HU, Yu Z, Schwartz O. A long-term study of 370 autotransplanted
6 premolars. Part III. Periodontal healing subsequent to transplantation. European Journal of
7 Orthodontics 1990;12:25-37.
8
9 Andreasen JO, Paulsen HU, Yu Z, Bayer T. A long-term study of 370 autotransplanted premolars.
10 Part IV. Root development subsequent to transplantation. European Journal of Orthodontics
11 1990;12:38-50.
12
13 Baumrind S, Korn EL, Isaacson RJ, West EE, Molthen R. Quantitative analysis of the orthodontic
14 and orthopedic effects of maxillary traction. American Journal of Orthodontics 1983;84:384-98.
15
16 Baumrind S, Korn EL, West EE. Prediction of mandibular rotation: an empirical test of clinician
17 performance. American Journal of Orthodontics 1984;86:371-85.
18
19 Baumrind S, Molthen R, West EE, Miller DM. Mandibular plane changes during maxillary
20 retraction. American Journal of Orthodontics 1978;74:32-40.
21
22 Baumrind S, Molthen R, West EE, Miller DM. Mandibular plane changes during maxillary
23 retraction. Part 2. American Journal of Orthodontics 1978;74:603-20.
24
25 Bedoya MM, Park JH. A review of the diagnosis and management of impacted maxillary canines.
26 Journal of the American Dental Association 2009;140:1485-93.
27
28 Brennan MM, Gianelly AA. The use of the lingual arch in the mixed dentition to resolve incisor
29 crowding. American Journal of Orthodontics and Dentofacial Orthopedics 2000;117:81-5.
30
31 Broadbent BH Sr, Broadbent BH Jr, Golden WH. Bolton standards of dentofacial developmental
32 growth. St. Louis: Mosby; 1975.
33
34 Burstone CR. Deep overbite correction by intrusion. American Journal of Orthodontics 1977;72:1-
35 22.
36
37 Cevidanes L, Baccetti T, Franchi L, McNamara JA Jr, De Clerck H. Comparison of two protocols
38 for maxillary protraction: bone anchors versus face mask with rapid maxillary expansion. Angle
39 Orthodontist 2010;80:799-806.
40
41 Chang CA, Fields HW, Beck FM, Springer NC, Firestone AR, Rosenstiel S, Christensen JC. Smile
42 esthetics from patients' perspective for faces of varying attractiveness. American Journal of
43 Orthodontics and Dentofacial Orthopedics 2011;140:e171-80.
44
45 Cordasco G, Matarese G, Rustico L, Fastuca S, Caprioglio A, Lindauer SJ, Nucera R. Efficacy of
46 orthopedic treatment with protraction facemask on skeletal Class III malocclusion: a systematic
47 review and meta-analysis. Orthodontics and Craniofacial Research 2014;17:133-43.
48
49 Czochrowska EM, Stenvik A, Bjercke B, Zachrisson BU. Outcome of tooth transplantation: survival
50 and success rates 17-41 years post treatment. American Journal of Orthodontics and Dentofacial
51 Orthopedics 2002;121:110-9.

1 Dale JG, Dale HC. Interceptive guidance of occlusion, with emphasis on diagnosis. In: Graber LW,
2 Vanarsdall RL, Vig KWL. Orthodontics: current principles and techniques. 5th ed. Philadelphia:
3 Elsevier/Mosby; 2012. p. 423-476.
4

5 Downs WB. Variations in facial relationships; their significance in treatment and prognosis.
6 American Journal of Orthodontics 1948;34:812-40.
7

8 English JD, Tadlock L, Briss BS, Litschel K. Diagnosis of orthodontic problems. In: English JD,
9 Akyalcin S, Peltomäki T, Litschel, K. Mosby's orthodontic review. 2nd ed. St. Louis: Elsevier Mosby,
10 2015. p. 60-97.
11

12 Fields HW. Complex nonskeletal problems in preadolescent children: preventive and interceptive
13 treatment. In: Proffit WR, Fields HW, Sarver DM, Ackerman, JL. Contemporary orthodontics. 5th
14 ed. St. Louis: Elsevier/Mosby; 2013. p. 446-471.
15

16 Fields HW, Proffit WR. Treatment of skeletal problems in children and preadolescents. In: Proffit
17 WR, Fields HW, Sarver DM, Ackerman, JL. Contemporary orthodontics. 5th ed. St. Louis:
18 Elsevier/Mosby; 2013. p. 472-528.
19

20 Foersch M, Jacobs C, Wriedt S, Hechtner M, Wehrbein H. Effectiveness of maxillary protraction
21 using facemask with or without maxillary expansion: a systematic review and meta-analysis.
22 Clinical Oral Investigations 2015;19:1181-92.
23

24 Franchi L, Contardo L, Primožič J, Perinetti G. Clinical alteration of mandibular growth: what we
25 know after 40 years. In: McNamara JA Jr, De Koster KY. The 40th Moyers Symposium: looking
26 back...looking forward. Ann Arbor: The University of Michigan; 2014. p. 263-285.
27

28 Ge YS, Liu J, Chen L, Han JL, Guo X. Dentofacial effects of two facemask therapies for maxillary
29 protraction. Angle Orthodontist 2012;82:1083-91.
30

31 Graber TM. Functional appliances. In: Graber TM, Vanarsdall RL, Vig KML. Orthodontics: current
32 principles and techniques. 4th ed. St. Louis: Elsevier/Mosby; 2005. p. 493-542.
33

34 Grec RH, Janson G, Branco NC, Moura-Grec PG, Patel MP, Castanha Henriques JF. Intraoral
35 distalizer effects with conventional and skeletal anchorage: a meta-analysis. American Journal of
36 Orthodontics and Dentofacial Orthopedics 2013;143:602-15.
37

38 Greenlee GM, Huang GJ, Chen SS, Chen J, Koepsell T, Hujoel P. Stability of treatment for
39 anterior open-bite malocclusion: a meta-analysis. American Journal of Orthodontics and
40 Dentofacial Orthopedics 2011;139:154-69.
41

42 Gu Y. Factors contributing to stability of protraction facemask treatment of Class III malocclusion.
43 Australian Orthodontic Journal 2010;26:171-7.
44

45 Hartsfield JK. Genetics and orthodontics. In: Graber LW, Vanarsdall RL, Vig KWL. Orthodontics:
46 current principles and techniques. 5th ed. Philadelphia: Elsevier/Mosby; 2012. p. 139-156.
47

48 Harvold EP. The activator in interceptive orthodontics. St. Louis: CV Mosby; 1974.
49

1 Haryett RD, Hansen FC, Davidson PO, Sandilands ML. Chronic thumb-sucking: the psychologic
2 effects and the relative effectiveness of various methods of treatment. *American Journal of*
3 *Orthodontics* 1967;53:569-85.
4
5 Husain J, Burden D, McSherry P, Morris D, Allen M, Clinical Standards Committee of the Faculty
6 of Dental Surgery, Royal College of Surgeons of England. National clinical guidelines for
7 management of the palatally ectopic maxillary canine. *British Dental Journal* 2012;213:171-6.
8
9 Huynh NT, Desplats E, Almeida FR. Orthodontics treatments for managing obstructive sleep
10 apnea syndrome in children: a systematic review and meta-analysis. *Sleep Medicine Reviews*
11 2016; 25:84-94.
12
13 Jacobson A. The "Wits" appraisal of jaw disharmony. *American Journal of Orthodontics*
14 1975;67:125-38.
15
16 Janson G, Sathler R, Fernandes TM, Branco NC, Freitas MR. Correction of Class II malocclusion
17 with Class II elastics: a systematic review. *American Journal of Orthodontics and Dentofacial*
18 *Orthopedics* 2013;143:383-92.
19
20 Kennedy DB, Turley PK. The clinical management of ectopically erupting first permanent molars.
21 *American Journal of Orthodontics and Dentofacial Orthopedics* 1987;92:336-45.
22
23 Ker AJ, Chan R, Fields HW, Beck M, Rosenstiel S. Esthetic and smile characteristics from the
24 layperson's perspective: a computer-based survey study. *Journal of the American Dental*
25 *Association* 2008;139:1318-27.
26
27 Kim YH, Vietas JJ. Anteroposterior dysplasia indicator: an adjunct to cephalometric differential
28 diagnosis. *American Journal of Orthodontics* 1978;73:619-33.
29
30 Kokich VO Jr, Kiyak HA, Shapiro PA. Comparing the perception of dentists and lay people to
31 altered dental esthetics. *Journal of Esthetic Dentistry* 1999;11:311-24.
32
33 Martina R, Cioffi I, Farella M, Leone P, Manzo P, Matarese G, Portelli M, Nucera R, Cordasco G.
34 Transverse changes determined by rapid and slow maxillary expansion--a low-dose CT-based
35 randomized controlled trial. *Orthodontics and Craniofacial Research* 2012;15:159-68.
36
37 McNamara JA Jr. A method of cephalometric evaluation. *American Journal of Orthodontics*
38 1984;86:449-69.
39
40 McNamara JA, McNamara L, Graber LW. Optimizing orthodontic and dentofacial orthopedic
41 treatment timing. In: Graber LW, Vanarsdall RL, Vig KWL. *Orthodontics: current principles and*
42 *techniques*. 5th ed. Philadelphia: Elsevier/Mosby; 2012. p. 477-514.
43
44 Millett DT, Cunningham SJ, O'Brien KD, Benson P, Williams A, de Oliveira CM. Orthodontic
45 treatment for deep bite and retroclined upper front teeth in children. *Cochrane Database*
46 *Systematic Reviews* 2006 Oct 18;(4):CD005972.
47
48 Nazarali N, Altalibi M, Nazarali S, Major MP, Flores-Mir C, Major PW. Mandibular advancement
49 appliances for the treatment of paediatric obstructive sleep apnea: a systematic review. *European*
50 *Journal of Orthodontics* 2015;37:618-26.
51

1 O'Brien K, Wright J, Conboy F, Sanjie Y, Mandall N, Chadwick S, Connolly I, Cook P, Birnie D,
2 Hammond M, Harradine N, Lewis D, McDade C, Mitchell L, Murray A, O'Neill J, Read M, Robinson
3 S, Roberts-Harry D, Sandler J, Shaw I. Effectiveness of early orthodontic treatment with the Twin-
4 block appliance: a multicenter, randomized, controlled trial. Part 1: Dental and skeletal effects.
5 American Journal of Orthodontics and Dentofacial Orthopedics 2003;124:234-43.
6
7 Okeson JP. Management of temporomandibular disorders and occlusion. 7th ed. St. Louis:
8 Elsevier/Mosby, 2013.
9
10 Opdebeeck H, Bell WH. The short face syndrome. American Journal of Orthodontics 1978;73:499-
11 511.
12
13 Perinetti G, Primožič J, Furlani G, Franchi L, Contardo L. Treatment effects of fixed functional
14 appliances alone or in combination with multibracket appliances: a systematic review and meta-
15 analysis. Angle Orthodontist 2015;85:480-92.
16
17 Proffit WR, Fields HW Jr, Moray LJ. Prevalence of malocclusion and orthodontic treatment need in
18 the United States: estimates from the NHANES III survey. International Journal of Adult
19 Orthodontics and Orthognathic Surgery 1998;13:97-106.
20
21 Proffit WR, Fields HW, Sarver DM. Orthodontic treatment planning: from problem list to specific
22 plan. In: Proffit WR, Fields HW, Sarver DM, Ackerman, JL. Contemporary orthodontics. 5th ed. St.
23 Louis: Elsevier/Mosby; 2013. p. 220-275.
24
25 Proffit WR, Sarver DM, Ackerman JL. Orthodontic diagnosis: the problem-oriented approach. In:
26 Proffit WR, Fields HW, Sarver DM, Ackerman, JL. Contemporary orthodontics. 5th ed. St. Louis:
27 Elsevier/Mosby; 2013. p. 150-219.
28
29 Proffit WR, Sarver DM. Special considerations in treatment for adults. In: Proffit WR, Fields HW,
30 Sarver DM, Ackerman, JL. Contemporary orthodontics. 5th ed. St. Louis: Elsevier/Mosby; 2013. p.
31 623-684.
32
33 Richardson ER. Racial differences in dimensional traits of the human face. Angle Orthodontist
34 1980;50:301-11.
35
36 Riedel RA. An analysis of dentofacial relationships. American Journal of Orthodontics
37 1957;43:103-19.
38
39 Saffer F, Lubianca Neto JF, Rösing C, Dias C, Closs L. Predictors of success in the treatment of
40 obstructive sleep apnea syndrome with mandibular repositioning appliance: a systematic review.
41 International Archives of Otorhinolaryngology 2015;19:80-5.
42
43 Sagne S, Thilander B. Transalveolar transplantation of maxillary canines. A critical evaluation of a
44 clinical procedure. Acta Odontologica Scandinavica 1997;55:1-8.
45
46 Sandler J, Murray A, Thiruvengkatachari B, Gutierrez R Speight P, O'Brien K. Effectiveness of 3
47 methods of anchorage reinforcement for maximum anchorage in adolescents: a 3-arm multicenter
48 randomized clinical trial. American Journal of Orthodontics and Dentofacial Orthopedics
49 2014;146:10-20.
50

1 Sarver DM, Yanosky M. Special considerations in diagnosis and treatment planning. In: Graber
2 LW, Vanarsdall RL, Vig KWL. Orthodontics: current principles and techniques. 5th ed.
3 Philadelphia: Elsevier/Mosby; 2012. p. 59-98.
4

5 Schendel SA, Eisenfeld J, Bell WH, Epker B, Mishelevich DJ. The long face syndrome: vertical
6 maxillary excess. American Journal of Orthodontics 1976;70:398-408.
7

8 Seehra J, Fleming PS, Mandall N, Dibiase AT. A comparison of two different techniques for early
9 correction of Class III malocclusion. Angle Orthodontist 2012;82:96-101.
10

11 Shaw WC. The influence of children's dentofacial appearance on their social attractiveness as
12 judged by peers and lay adults. American Journal of Orthodontics 1981;79:399-415.
13

14 Steiner CC. Cephalometries in clinical practice. Angle Orthodontist 1959;29:8-29.
15

16 Sugawara J, Mitani H. Facial growth of skeletal Class III malocclusion and the effects, limitations,
17 and long-term dentofacial adaptations to chin cap therapy. Seminars in Orthodontics 1997;3:244-
18 54.
19

20 Thiruvenkatachari B, Harrison JE, Worthington HV, O'Brien KD. Orthodontic treatment for
21 prominent upper front teeth (Class II malocclusion) in children. The Cochrane Database of
22 Systematic Reviews 2013;11:CD003452.
23

24 Tulloch JF, Proffit WR, Phillips C. Outcomes in a 2-phase randomized clinical trial of early Class II
25 treatment. American Journal of Orthodontics and Dentofacial Orthopedics 2004;125:657-67.
26

27 Ubaldo ED, Greenlee GM, Moore J, Sommers E, Bollen AM. Cephalometric analysis and long-
28 term outcomes of orthognathic surgical treatment for obstructive sleep apnoea. International
29 Journal of Oral and Maxillofacial Surgery 2015;44:752-9.
30

31 Vanarsdall RL, Musich DR. Adult interdisciplinary therapy: diagnosis and treatment. In: Graber
32 LW, Vanarsdall RL, Vig KWL. Orthodontics: current principles and techniques. 5th ed.
33 Philadelphia: Elsevier/Mosby; 2012. p. 843-96.
34

35 Westwood PV, McNamara JA Jr, Baccetti T, Franchi L, Sarver DM. Long-term effects of Class III
36 treatment with rapid maxillary expansion and facemask therapy followed by fixed appliances.
37 American Journal of Orthodontics and Dentofacial Orthopedics 2003;123:306-20.
38

39 Wheeler TT, McGorray SP, Dolce C, Taylor MG, King GJ. Effectiveness of early treatment of
40 Class II malocclusion. American Journal of Orthodontics and Dentofacial Orthopedics 2002;121:9-
41 17.
42

43 Wieslander L. Intensive treatment of severe Class II malocclusions with headgear-Herbst
44 appliance in the early mixed dentition. American Journal of Orthodontics 1984;86:1-13.
45

46 Wieslander L, Lagerström L. The effect of activator treatment on class II malocclusions. American
47 Journal of Orthodontics 1979;75:20-6.
48

49 Will L. Psychological aspects of orthodontics. In: Graber LW, Vanarsdall RL, Vig KWL.
50 Orthodontics: current principles and techniques. 5th ed. Philadelphia: Elsevier/Mosby; 2012. p. 99-
51 107.

1 Zachrisson BU, Buyukyilmaz T. Bonding in orthodontics. In: Graber LW, Vanarsdall RL, Vig KWL.
2 Orthodontics: current principles and techniques. 5th ed. Philadelphia: Elsevier/Mosby; 2012. p.
3 727-84.
4

5 Zhou Y, Long H, Ye N, Xue J, Yang X, Liao L, Lai W. The effectiveness of non-surgical maxillary
6 expansion: a meta-analysis. *European Journal of Orthodontics* 2014;36:233-42.
7

8 **Diagnostic and Treatment Considerations for Anomalies of Jaw Size, Relationship of Jaw**
9 **to Cranial Base, Dental Arch Relationship and Dental Alveolus**

10

11 Ackerman JL, Nguyen T, Proffit WR. The decision making process in orthodontics. In: Graber LW,
12 Vanarsdall RL, Vig KWL. Orthodontics: current principles and techniques. 5th ed. Philadelphia:
13 Elsevier/Mosby; 2012. p. 3-58.
14

15 Alexander RG, Sinclair PM, Goates LJ. Differential diagnosis and treatment planning for the adult
16 nonsurgical orthodontic patient. *American Journal of Orthodontics* 1986;89:95-112.
17

18 Arvystas MG. Treatment of anterior skeletal open-bite deformity. *American Journal of Orthodontics*
19 1977;72:147-64.
20

21 Bell WH, Jacobs JD, Legan HL. Treatment of Class II deep bite by orthodontic and surgical
22 means. *American Journal of Orthodontics* 1984;85:1-20.
23

24 Burstone CR. Deep overbite correction by intrusion. *American Journal of Orthodontics* 1977;72:1-
25 22.
26

27 Cangialosi TJ, Meistrell ME Jr, Leung MA, Ko JY. A cephalometric appraisal of edgewise Class II
28 nonextraction treatment with extraoral force. *American Journal of Orthodontics and Dentofacial*
29 *Orthopedics* 1988;93:315-24.
30

31 Carlotti AE, George R. Differential diagnosis and treatment planning of the surgical orthodontic
32 class III malocclusion. *American Journal of Orthodontics* 1981;79:424-36.
33

34 Chaconas SJ, de Alba y Levy JA. Orthopedic and orthodontic applications of the quad-helix
35 appliance. *American Journal of Orthodontics* 1977;72:422-8.
36

37 Cozza P, Baccetti T, Franchi L, De Toffol L, McNamara JA Jr. Mandibular changes produced by
38 functional appliances in Class II malocclusion: a systematic review. *American Journal of*
39 *Orthodontics and Dentofacial Orthopedics* 2006;129:599.e1-12; discussion e1-6.
40

41 D'Antò V, Bucci R, Franchi L, Rongo R, Michelotti A, Martina R. Class II functional orthopaedic
42 treatment: a systematic review of systematic reviews. *Journal of Oral Rehabilitation* 2015;42:624-
43 42.
44

45 El-Dawlatly MM, Fayed MM, Mostafa YA. Deep overbite malocclusion: analysis of the underlying
46 components. *American Journal of Orthodontics and Dentofacial Orthopedics* 2012;142:473-80.
47

48 Epker BN, Fish L. Surgical-orthodontic correction of open-bite deformity. *American Journal of*
49 *Orthodontics* 1977;71:278-99.
50

1 Epker BN, Wolford LM, Fish LC. Mandibular deficiency syndrome II. Surgical considerations for
2 mandibular advancement. *Oral Surgery, Oral Medicine, and Oral Pathology* 1978;45:349-63.
3
4 Frankel R, Frankel C. A functional approach to treatment of skeletal open bite. *American Journal*
5 *of Orthodontics* 1983;84:54-68.
6
7 Freeman CS, McNamara JA Jr, Baccetti T, Franchi L, Graff TW. Treatment effects of the bionator
8 and high-pull facebow combination followed by fixed appliances in patients with increased vertical
9 dimensions. *American Journal of Orthodontics and Dentofacial Orthopedics* 2007;131:184-95.
10
11 Gianelly AA, Arena SA, Bernstein L. A comparison of Class II treatment changes noted with the
12 light wire, edgewise, and Frankel appliances. *American Journal of Orthodontics* 1984;86:269-76.
13
14 Glassman AS, Nahigian SJ, Medway JM, Aronowitz HI. Conservative surgical orthodontic adult
15 rapid palatal expansion: sixteen cases. *American Journal of Orthodontics* 1984;86:207-13.
16
17 Glenn G, Sinclair PM, Alexander RG. Nonextraction orthodontic therapy: posttreatment dental and
18 skeletal stability. *American Journal of Orthodontics and Dentofacial Orthopedics* 1987;92:321-8.
19
20 Graber LW. Chin cup therapy for mandibular prognathism. *American Journal of Orthodontics*
21 1977;72:23-41.
22
23 Holdaway RA. A soft-tissue cephalometric analysis and its use in orthodontic treatment planning.
24 Part II. *American Journal of Orthodontics* 1984;85:279-93.
25
26 Jacobson A. The "Wits" appraisal of jaw disharmony. *American Journal of Orthodontics*
27 1975;67:125-38.
28
29 Kokich VG. Adult orthodontics in the 21st century: guidelines for achieving successful results.
30 *World Journal of Orthodontics* 2005;6 Suppl:14-23.
31
32 Liu S, Xu T, Zou W. Effects of rapid maxillary expansion on the midpalatal suture: a systematic
33 review. *European Journal of Orthodontics* 2015;37:651-5.
34
35 Magness WB. The mini-visualized treatment objective. *American Journal of Orthodontics and*
36 *Dentofacial Orthopedics* 1987;91:361-74.
37
38 Marşan G. Effects of activator and high-pull headgear combination therapy: skeletal,
39 dentoalveolar, and soft tissue profile changes. *European Journal of Orthodontics* 2007;29:140-8.
40
41 Moyers RE, Bookstein, FL, Hunter, WS. Section II: Diagnosis. In: Moyers RE. *Handbook of*
42 *orthodontics*. 4th ed. Chicago: Year Book Medical Publishers; 1988. p. 165-301.
43
44 McNamara JA Jr, Hulse SA. The Frankel appliance (FR2): model preparation and appliance
45 construction. *American Journal of Orthodontics* 1981;80:478-95.
46
47 McNamara JA Jr. An orthopedic approach to the treatment of Class III malocclusion in young
48 patients. *Journal of Clinical Orthodontics* 1987;21:598-608.
49

1 McNamara JA, McNamara L, Graber LW. Optimizing orthodontic and dentofacial orthopedic
2 treatment timing. In: Graber LW, Vanarsdall RL, Vig KWL. Orthodontics: current principles and
3 techniques. 5th ed. Philadelphia: Elsevier/Mosby; 2012. p. 477-514.
4

5 Nahoum HI. Vertical proportions: a guide for prognosis and treatment in anterior open-bite.
6 American Journal of Orthodontics 1977;72:128-46.
7

8 Pancherz H. A cephalometric analysis of skeletal and dental changes contributing to Class II
9 correction in activator treatment. American Journal of Orthodontics 1984;85:125-34.
10

11 Pearson LE. Vertical control in fully-banded orthodontic treatment. Angle Orthodontist
12 1986;56:205-24.
13

14 Pearson LE. Treatment of a severe openbite excessive vertical pattern with an eclectic non-
15 surgical approach. Angle Orthodontist 1991;61:71-6.
16

17 Pfeiffer JP, Grobety D. A philosophy of combined orthopedic-orthodontic treatment. American
18 Journal of Orthodontics 1982;81:185-201.
19

20 Poulton DR, Ware WH. Increase in mandibular and chin projection with orthognathic surgery.
21 American Journal of Orthodontics 1985;87:363-76.
22

23 Proffit WR, White RP Jr. Combined surgical-orthodontic treatment: how did it evolve and what are
24 the best practices now? American Journal of Orthodontics and Dentofacial Orthopedics
25 2015;147:S205-15.
26

27 Richardson ER. Racial differences in dimensional traits of the human face. Angle Orthodontist
28 1980;50:301-11.
29

30 Sakamoto T. Effective timing for the application of orthopedic force in the skeletal class III
31 malocclusion. American Journal of Orthodontics 1981;80:411-6.
32

33 Skieller V, Bjork A, Linde-Hansen T. Prediction of mandibular growth rotation evaluated from a
34 longitudinal implant sample. American Journal of Orthodontics 1984;86:359-70.
35

36 Solano-Hernández B, Antonarakis GS, Scolozzi P, Kiliaridis S. Combined orthodontic and
37 orthognathic surgical treatment for the correction of skeletal anterior open-bite malocclusion: a
38 systematic review on vertical stability. Journal of Oral and Maxillofacial Surgery 2013;71:98-109.
39

40 Turpin DL. Befriend your oral and maxillofacial radiologist. American Journal of Orthodontics and
41 Dentofacial Orthopedics 2007;131:697.
42

43 Wendell PD, Nanda R, Sakamoto T, Nakamura, S. The effects of chin cup therapy on the
44 mandible: a longitudinal study. American Journal of Orthodontics 1985;87:265-74.
45

46 Wieslander L. Intensive treatment of severe Class II malocclusions with headgear-Herbst
47 appliance in the early mixed dentition. American Journal of Orthodontics 1984;86:1-13.
48

49 Williams S, Andersen CE. The morphology of the potential Class III skeletal pattern in the growing
50 child. American Journal of Orthodontics 1986;89:302-11.
51

1 Zuccati G, Casci S, Doldo T, Clauser C. Expansion of maxillary arches with crossbite: a
2 systematic review of RCTs in the last 12 years. *European Journal of Orthodontics* 2013;35:29-37.
3

4 **Diagnostic and Treatment Considerations for Anomalies of Tooth Position, Discrepancies**
5 **of Tooth Size, Arch Length and Arch Form**
6

7 Bolton WA. The clinical application of a tooth-size analysis. *American Journal Orthodontics*
8 1962;48:504-29.
9

10 Clark JD, Williams JK. The management of spacing in the maxillary incisor region. *British Journal*
11 *of Orthodontics* 1978;5:35-9.
12

13 Dale JG, Dale HC. Interceptive guidance of occlusion, with emphasis on diagnosis. In: Graber LW,
14 Vanarsdall RL, Vig KWL. *Orthodontics: current principles and techniques*. 5th ed. Philadelphia:
15 Elsevier/Mosby; 2012. p. 423-476.
16

17 Dewel BF. Serial extraction in orthodontics: indications, objectives, and treatment procedures.
18 *American Journal of Orthodontics* 1954;40:906-26.
19

20 Feldmann I, Bondemark L. Orthodontic anchorage: a systematic review. *Angle Orthodontist*
21 2006;76:493-501.
22

23 Howe RP, McNamara JA Jr, O'Connor KA. An examination of dental crowding and its relationship
24 to tooth size and arch dimension. *American Journal of Orthodontics* 1983;83:363-73.
25

26 Jerrold L, Lowenstein LJ. The midline: diagnosis and treatment. *American Journal of Orthodontics*
27 *and Dentofacial Orthopedics* 1990;97:453-62.
28

29 Lopes Filho H, Maia LH, Lau TC, de Souza MM, Maia LC. Early vs late orthodontic treatment of
30 tooth crowding by first premolar extraction: a systematic review. *Angle Orthodontist* 2015;85:510-
31 7.
32

33 Lundstrom A. The aetiology of crowding of the teeth (based on studies of twins and on
34 morphological investigations) and its bearing on orthodontic treatment (expansion or extraction).
35 *European Orthodontic Society Transactions* 1951;176-91.
36

37 McKeown M. The diagnosis of incipient arch crowding in children. *New Zealand Dental Journal*
38 1981;77:93-96.
39

40 Mills LF. Arch width, arch length, and tooth size in young adult males. *Angle Orthodontist*
41 1964;34:124-9.
42

43 Mills LF. Epidemiologic studies of occlusion. IV. The prevalence of malocclusion in a population of
44 1,455 school children. *Journal of Dental Research* 1966;45:332-6.
45

46 Moorrees CF, Reed RB. Biometrics of crowding and spacing of the teeth in the mandible.
47 *American Journal of Physical Anthropology* 1954;12:77-88.
48

49 Moyers RE. Standards of human occlusal development. Craniofacial growth series No. 5 Ann
50 Arbor, MI: Center for Human Growth and Development, University of Michigan; 1976.
51

1 O'Shaughnessy KW, Koroluk LD, Phillips C, Kennedy DB. Efficiency of serial extraction and late
2 premolar extraction cases treated with fixed appliances. *American Journal of Orthodontics and*
3 *Dentofacial Orthopedics* 2011;139:510-6.
4
5 Peck H, Peck S. An index for assessing tooth shape deviations as applied to the mandibular
6 incisors. *American Journal of Orthodontics* 1972;61:384-401.
7
8 Poulton DR. The influence of extraoral traction. *American Journal of Orthodontics* 1967;53:8-18.
9
10 Proffit WR, Fields HW, Sarver DM. Orthodontic treatment planning: from problem list to specific
11 plan. In: Proffit WR, Fields HW, Sarver DM, Ackerman, JL. *Contemporary orthodontics*. 5th ed. St.
12 Louis: Elsevier/Mosby; 2013. p. 220-275.
13
14 Proffit WR, Sarver DM, Ackerman JL. Orthodontic diagnosis: the problem-oriented approach. In:
15 Proffit WR, Fields HW, Sarver DM, Ackerman, JL. *Contemporary orthodontics*. 5th ed. St. Louis:
16 Elsevier/Mosby; 2013. p. 150-219.
17
18 Sheridan JJ. Air-rotor stripping update. *Journal of Clinical Orthodontics* 1987;21:781-8.
19
20 Terwilliger KF. Treatment in the mixed dentition. *Angle Orthodontist* 1950;20:109-13.
21
22 Vaden JL, Klontz HA, Dale JG. Standard Edgewise: Tweed-Merrifield philosophy, diagnosis,
23 treatment planning and force systems. In: Graber LW, Vanarsdall RL, Vig KWL. *Orthodontics:*
24 *current principles and techniques*. 5th ed. Philadelphia: Elsevier/Mosby; 2012. p. 517-59.
25
26 **Diagnostic and Treatment Considerations for Abnormalities of the Dentition (number, size**
27 **and shape), Vitality, Eruption Pattern and Periodontal Support**
28
29 Andreasen JO, Paulsen HU, Yu Z, Bayer T, Schwartz O. A long-term study of 370
30 autotransplanted premolars. Part II. Tooth survival and pulp healing subsequent to transplantation.
31 *European Journal of Orthodontics* 1990;12:14-24.
32
33 Andrade DC, Loureiro CA, Araujo VE, Riera R, Atallah AN. Treatment for agenesis of maxillary
34 lateral incisors: a systematic review. *Orthodontics and Craniofacial Research* 2013;16:129-36.
35
36 Becker A, Chaushu S, Casap-Caspi N. Cone-beam computed tomography and the orthosurgical
37 management of impacted teeth. *Journal of the American Dental Association* 2010;141 Suppl
38 3:14S-8S.
39
40 Brook AH, Jernvall J, Smith RN, Hughes TE, Townsend GC. The dentition: the outcomes of
41 morphogenesis leading to variations of tooth number, size and shape. *Australian Dental Journal*
42 2014;59 Suppl 1:131-42.
43
44 Carter K, Worthington S. Morphologic and demographic predictors of third molar agenesis: a
45 systematic review and meta-analysis. *Journal of Dental Research* 2015;94:886-94.
46
47 Chaushu S, Kaczor-Urbanowicz K, Zadurska M, Becker A. Predisposing factors for severe incisor
48 root resorption associated with impacted maxillary canines. *American Journal of Orthodontics and*
49 *Dentofacial Orthopedics* 2015;147:52-60.
50

1 Chaushu S, Becker T, Becker A. Impacted central incisors: factors affecting prognosis and
2 treatment duration. *American Journal of Orthodontics and Dentofacial Orthopedics* 2015;147:355-
3 62.
4
5 Dibase D. Mucous membrane and delayed eruption. *Transactions of the British Society for the*
6 *Study of Orthodontics* 1969/70;56:149-58.
7
8 Ericson S, Kurol J. Early treatment of palatally erupting maxillary canines by extraction of the
9 primary canines. *European Journal of Orthodontics* 1988;10:283-95.
10
11 Hua F, He H, Ngan P, Bouzid W. Prevalence of peg-shaped maxillary permanent lateral incisors: a
12 meta-analysis. *American Journal of Orthodontics and Dentofacial Orthopedics* 2013;144:97-109.
13
14 Joondeph DR, McNeill RW. Congenitally absent second premolars: an interceptive approach.
15 *American Journal of Orthodontics* 1971;59:50-66.
16
17 Kerekes-Máthé B, Brook AH, Mártha K, Székely M, Smith RN. Mild hypodontia is associated with
18 smaller tooth dimensions and cusp numbers than in controls. *Archives of Oral Biology*
19 2015;60:1442-9.
20
21 Kurol J, Thilander B. Infraocclusion of primary molars and the effect on occlusal development, a
22 longitudinal study. *European Journal of Orthodontics* 1984;6:277-93.
23
24 Luu NS, Mandich MA, Tieu LD, Kaipatur N, Flores-Mir C. The validity and reliability of mixed-
25 dentition analysis methods: a systematic review. *Journal of the American Dental Association*
26 2011;142:1143-53.
27
28 Mitchell L, Bennett TG. Supernumerary teeth causing delayed eruption-a retrospective study.
29 *British Journal of Orthodontics* 1992;19:41-6.
30
31 Moorrees CFA. The dentition of the growing child; a longitudinal study of dental
32 development between 3 and 18 years of age. Cambridge: Harvard University Press; 1959.
33
34 Naoumova J, Kurol J, Kjellberg H. A systematic review of the interceptive treatment of palatally
35 displaced maxillary canines. *European Journal of Orthodontics* 2011;33:143-9.
36
37 Odeh R, Mihailidis S, Townsend G, Lähdesmäki R, Hughes T, Brook A. Prevalence of
38 infraocclusion of primary molars determined using a new 2D image analysis methodology.
39 *Australian Dental Journal* 2015 Jul 14 [Epub ahead of print].
40
41 Odeh R, Townsend G, Mihailidis S, Lähdesmäki R, Hughes T, Brook A. Infraocclusion: dental
42 development and associated dental variations in singletons and twins. *Archives of Oral Biology*
43 2015;60:1394-402.
44
45 Parkin N, Benson PE, Thind B, Shah A. Open versus closed surgical exposure of canine teeth that
46 are displaced in the roof of the mouth. *Cochrane Database of Systematic Reviews*
47 2008;(4):CD006966.
48
49 Paulsen HU, Andreasen JO, Schwartz O. Pulp and periodontal healing, root development and root
50 resorption subsequent to transplantation and orthodontic rotation: a long term study of

1 autotransplanted premolars. American Journal of Orthodontics and Dentofacial Orthopedics
2 1995;108:630-40.
3
4 Paulsson L, Bondemark L, Soderfeldt B. A systematic review of the consequences of premature
5 birth on palatal morphology, dental occlusion, tooth-crown dimensions, and tooth maturity and
6 eruption. Angle Orthodontist 2004;74:269-79.
7
8 Peck S, Peck L. Classification of maxillary tooth transpositions. American Journal of Orthodontics
9 and Dentofacial Orthopedics 1995;107:505-17.
10
11 Poelmans S, Clijmans M, Fieuws S, Willems G. Cephalometric appraisal of post-treatment tooth
12 eruption: a 20 year follow-up study. European Journal of Orthodontics 2015 Mar 28 [Epub ahead
13 of print].
14
15 Pulver P. The etiology and prevalence of ectopic eruption of the maxillary first permanent molar.
16 ASDC Journal of Dentistry for Children 1968;35:138-46.
17
18 Rakhshan V, Rakhshan H. Meta-analysis and systematic review of the number of non-syndromic
19 congenitally missing permanent teeth per affected individual and its influencing factors. European
20 Journal of Orthodontics 2016;38:170-7.
21
22 Schatz JP, Joho JP. Indications of autotransplantation of teeth in orthodontic problem cases.
23 American Journal of Orthodontics and Dentofacial Orthopedics 1994;106:351-7.
24
25 Thilander B. Orthodontic space closure versus implant placement in subjects with missing teeth.
26 Journal of Oral Rehabilitation 2008;35 Suppl 1:64-71.
27
28 **Diagnostic and Treatment Considerations for Dentofacial Functional Abnormalities**
29
30 American Association of Orthodontists. Glossary of dentofacial orthopedic terms. St. Louis:
31 American Association of Orthodontists; 2012.
32
33 Andrianopoulos MV, Hanson ML. Tongue-thrust and the stability of overjet correction. Angle
34 Orthodontist 1987;57:121-35.
35
36 Baumrind S, Korn EL, Isaacson RJ, West EE, Molthen R. Superimpositional assessment of
37 treatment-associated changes in the temporomandibular joint and the mandibular symphysis.
38 American Journal of Orthodontics 1983;84:443-65.
39
40 Behrents RG, White RA. TMJ research: responsibility and risk. American Journal of Orthodontics
41 and Dentofacial Orthopedics 1992;101:1-3.
42
43 Burton MJ, Glasziou PP, Chong LY, Venekamp RP. Tonsillectomy or adenotonsillectomy versus
44 non-surgical treatment for chronic/recurrent acute tonsillitis. Cochrane Database of Systematic
45 Reviews 2014;11:CD001802.
46
47 Dibbets JM, van der Weele LT. The prevalence of joint noises as related to age and gender.
48 Journal of Craniomandibular Disorders 1992;6:157-60.
49

1 Fields HW, Warren DW, Black K, Phillips CL. Relationship between vertical dentofacial
2 morphology and respiration in adolescents. *American Journal of Orthodontics and Dentofacial*
3 *Orthopedics* 1991;99:147-54.
4
5 Graber TM. The "three M's": Muscles, malformation, and malocclusion. *American Journal*
6 *Orthodontics* 1963;49:418-50.
7
8 Graber TM. Postmortems in posttreatment adjustment. *American Journal of Orthodontics*
9 1966;52:331-52.
10
11 Grummons D. *Orthodontics for the TMJ-TMD patient*. Scottsdale, Ariz.: Wright, & Co. Publishers;
12 1994.
13
14 Harvold EP, Tomer BS, Vargervik K, Chierici G. Primate experiments on oral respiration. *American*
15 *Journal of Orthodontics* 1981;79:359-72.
16
17 Haryett RD, Hansen FC, Davidson PO. Chronic thumb sucking. A second report on treatment and
18 its psychological effects. *American Journal of Orthodontics* 1970;57:164-78.
19
20 Ingervall B. Orthodontic treatment in adults with temporomandibular dysfunction symptoms.
21 *American Journal of Orthodontics* 1978;73:551-9.
22
23 Jones AG, Bhatia S. A study of nasal respiratory resistance and craniofacial dimensions in white
24 and West Indian black children. *American Journal of Orthodontics and Dentofacial Orthopedics*
25 1994;106:34-9.
26
27 Katyal V, Pamula Y, Martin AJ, Daynes CN, Kennedy JD, Sampson WJ. Craniofacial and upper
28 airway morphology in pediatric sleep-disordered breathing: systematic review and meta-analysis.
29 *American Journal of Orthodontics and Dentofacial Orthopedics* 2013;143:20-30.
30
31 Kerr WJ, McWilliam JS, Linder-Aronson S. Mandibular form and position related to changed mode
32 of breathing- a five-year longitudinal study. *Angle Orthodontist* 1989;59:91-6.
33
34 Larsson EF, Dahlin KG. The prevalence and the etiology of the initial dummy- and finger-sucking
35 habit. *American Journal of Orthodontics* 1985;87:432-5.
36
37 Linder-Aronson S. Adenoids. Their effect on mode of breathing and nasal airflow and their
38 relationship to characteristics of the facial skeleton and the dentition. A biometric, rhino-manometric
39 and cephalometro-radiographic study on children with and without adenoids. *Acta Oto-*
40 *Laryngologica. Supplement (Oslo)* 1970;265:1-132.
41
42 Linder-Aronson S, Leighton BC. A longitudinal study of the development of the posterior
43 nasopharyngeal wall between 3 and 16 years of age. *European Journal of Orthodontics*
44 1983;5:47-58.
45
46 Linder-Aronson S, Woodside DG, Hellsing E, Emerson W. Normalization of incisor position after
47 adenoidectomy. *American Journal of Orthodontics and Dentofacial Orthopedics* 1993;103:412-27.
48
49 Major MP, Saltaji H, El-Hakim H, Witmans M, Major P, Flores-Mir C. The accuracy of diagnostic
50 tests for adenoid hypertrophy: a systematic review. *Journal of the American Dental Association*
51 2014;145:247-54.

1 Mason RM. Orthodontic perspectives on orofacial myofunctional therapy. *International Journal of*
2 *Oral and Maxillofacial Surgery* 1988;14:49-55.
3
4 McNeill C. *Cranio-mandibular disorders: guidelines for evaluation, diagnosis, and management.*
5 Chicago: Quintessence Publishing Company; 1990.
6
7 Nagaiwa M, Gunjigake K, Yamaguchi K. The effect of mouth breathing on chewing efficiency.
8 *Angle Orthodontist* 2016;86:227-34.
9
10 Niinimaa V, Cole P, Mintz S, et al. Oronasal distribution of respiratory airflow. *Respiration*
11 *Physiology* 1981;43:69-75.
12
13 Proffit WR. Lingual pressure patterns in the transition from tongue thrust to adult swallowing.
14 *Archives of Oral Biology* 1972;17:555-63.
15
16 Roth RH. Functional occlusion for the orthodontist. Part III. *Journal of Clinical Orthodontics* 1981;
17 15:174-9, 182-98.
18
19 Sadowsky S, BeGole EA. Long-term status of temporomandibular joint function and functional
20 occlusion after orthodontic treatment. *American Journal of Orthodontics* 1980;78:201-12.
21
22 Stringert HG, Worms FW. Variations in skeletal and dental patterns in patients with structural and
23 functional alterations of the temporomandibular joint: a preliminary report. *American Journal of*
24 *Orthodontics* 1986; 89:285-97.
25
26 Subtelny JD. Oral habits - studies in form, function and therapy. *Angle Orthodontist* 1973;43:349-
27 83.
28
29 Tamari K, Murakami T, Takahama Y. The dimensions of the tongue in relation to its motility.
30 *American Journal of Orthodontics and Dentofacial Orthopedics* 1991;99:140-6.
31
32 te Veldhuis EC, te Veldhuis AH, Koudstaal MJ. Treatment management of children with juvenile
33 idiopathic arthritis with temporomandibular joint involvement: a systematic review. *Oral Surgery,*
34 *Oral Medicine, Oral Pathology and Oral Radiology* 2014;117:581-9.e2.
35
36 Valladares-Neto J, Cevidanes LH, Rocha WC, Almeida Gde A, Paiva JB, Rino-Neto J. TMJ
37 response to mandibular advancement surgery: an overview of risk factors. *Journal of Applied Oral*
38 *Science* 2014;22:2-14.
39
40 Vig KW. Orthodontic considerations applied to craniofacial dysmorphology. *Cleft Palate Journal*
41 1990;27:141-5.
42
43 Vig PS, Sarver DM, Hall DJ, Warren DW. Quantitative evaluation of nasal airflow in relation to
44 facial morphology. *American Journal of Orthodontics* 1981;79:263-72.
45
46 Vig PS, Showfety KJ, Phillips C. Experimental manipulation of head posture. *American Journal of*
47 *Orthodontics* 1980;77:258-68.
48
49 Wang XD, Zhang JN, Gan YH, Zhou YH. Current understanding of pathogenesis and treatment of
50 TMJ osteoarthritis. *Journal of Dental Research* 2015;94:666-73.
51

1 Watson RM Jr, Warren DW, Fischer ND. Nasal resistance, skeletal classification and mouth
2 breathing in orthodontic patients. *American Journal of Orthodontics* 1968;54:367-79.

3
4 Williamson EH. Temporomandibular dysfunction in pretreatment adolescent patients. *American*
5 *Journal of Orthodontics* 1977;72:429-33.

6 **Orthodontic Considerations for Craniofacial Anomalies, Cleft Lip and Palate**

7
8
9 American Cleft Palate-Craniofacial Association. Parameters for evaluation and treatment of
10 patients with cleft lip/palate or other craniofacial anomalies. *Cleft Palate Craniofacial Journal*
11 1993;30 Suppl:S1-16; current versions available on ACPA website: www.acpa-cpf.org.

12
13 Boyne PJ, Sands NR. Secondary bone grafting of residual alveolar and palatal clefts. *Journal of*
14 *Oral Surgery* 1972;30:87-92.

15
16 Bongaarts CA, Kuijpers-Jagtman AM, van't Hof MA, Prah-Andersen B. The effect of infant
17 orthopedics on the occlusion of the deciduous dentition in children with complete unilateral cleft lip
18 and palate (Dutchcleft). *Cleft Palate-Craniofacial Journal* 2004;41:633-41.

19
20 Cooper HK. Integration of services in the treatment of cleft lip and palate. *Journal of the American*
21 *Dental Association* 1953;47:27-32.

22
23 Daskalogiannakis J, Mercado A, Russell K, Hathaway R, Dugas G, Long RE Jr, Cohen M, Semb
24 G, Shaw W. The Americleft study: an inter-center study of treatment outcomes for patients with
25 unilateral cleft lip and palate part 3. Analysis of craniofacial form. *Cleft Palate-Craniofacial Journal*
26 2011;48:252-8.

27
28 Eskenazi LB, Schendel SA. An analysis of Le Fort I maxillary advancement in cleft lip and palate
29 patients. *Plastic and Reconstructive Surgery* 1992;90:779-86.

30
31 Gorlin RJ, Pindborg JJ. *Syndromes of the head and neck*. New York: McGraw-Hill; 1990.

32
33 Graber TM. Craniofacial morphology in cleft palate and cleft lip deformities. *Surgery, Gynecology*
34 *and Obstetrics* 1949;88:359-69.

35
36 Grayson BH, Santiago PE, Brecht LE, Cutting CB. Presurgical nasopalveolar molding in infants with
37 cleft lip and palate. *Cleft Palate-Craniofacial Journal* 1999;36:486-98.

38
39 Hathaway R, Daskalogiannakis J, Mercado A, Russell K, Long RE Jr, Cohen M, Semb G, Shaw
40 W. The Americleft study: an inter-center study of treatment outcomes for patients with unilateral
41 cleft lip and palate part 2. Dental arch relationships. *Cleft Palate-Craniofacial Journal* 2011;48:244-
42 51.

43
44 Kuijpers-Jagtman AM, Long RE Jr. State of the art - The influence of surgery and orthopedic
45 treatment on maxillofacial growth and maxillary arch development in patients treated for orofacial
46 clefts. *Cleft Palate-Craniofacial Journal* 2000;37:527e1-12.

47
48 Liou EJ, Tsai WC. A new protocol for maxillary protraction in cleft patients: repetitive weekly
49 protocol of alternate rapid maxillary expansions and constrictions. *Cleft Palate-Craniofacial Journal*
50 2005;42:121-7.

51

1 Long RE Jr, Semb G, Shaw WC. State of the art - Orthodontic treatment of the patient with
2 complete clefts of the lip, alveolus and palate. Lessons of the past 60 years. *Cleft Palate-*
3 *Craniofacial Journal* 2000;37:533e1-13.
4
5 Losee JE, Kirschner RE. *Comprehensive Cleft Care*. New York: McGraw-Hill Medical; 2009.
6
7 Mars M, Plint DA, Houston WJ, Bergland O, Semb G. The Goslon Yardstick: a new system of
8 assessing dental arch relationships in children with unilateral clefts of the lip and palate. *Cleft*
9 *Palate Journal* 1987;24:314-22.
10
11 Mercado AM, Vig KWL. Orthodontic principles in the management of oro-facial clefts. In: Losee
12 JE, Kirschner RE. *Comprehensive Cleft Care*. New York: McGraw-Hill Medical; 2009. p. 721-47.
13
14 Mølsted K, Brattström V, Prah-Andersen B, Shaw WC, Semb G. The Eurocleft study: intercenter
15 study of treatment outcome in patients with complete cleft lip and palate. Part 3: dental arch
16 relationships. *Cleft Palate-Craniofacial Journal* 2005;42:78-82.
17
18 Nollet PJ, Katsaros C, Van't Hof MA, Kuijpers-Jagtman AM. Treatment outcome in unilateral cleft
19 lip and palate evaluated with the GOSLON yardstick: a meta-analysis of 1236 patients. *Plastic and*
20 *Reconstructive Surgery* 2005;116:1255-62.
21
22 Polley JW, Figueroa AA. Rigid external distraction: its application in cleft maxillary deformities.
23 *Plastic and Reconstructive Surgery* 1998;102:1360-74.
24
25 Prah C, Kuijpers-Jagtman AM, van't Hof MA, Prah-Andersen B. A randomized prospective clinical
26 trial into the effect of infant orthopaedics on maxillary arch dimensions in unilateral cleft lip and
27 palate. *European Journal of Oral Sciences* 2001;109:297-305.
28
29 Ross RB. Treatment variables affecting facial growth in complete unilateral cleft lip and palate.
30 *Cleft Palate Journal* 1987;24:5-77.
31
32 Russell K, Long RE Jr, Hathaway R, Daskalogiannakis J, Mercado A, Cohen M, Semb G, Shaw
33 W. The Americleft study: an inter-center study of treatment outcomes for patients with unilateral
34 cleft lip and palate part 5. General discussion and conclusions. *Cleft Palate-Craniofacial Journal*
35 2011;48:265-70.
36
37 Semb G, Brattström V, Mølsted K, Prah-Andersen B, Shaw WC. The Eurocleft study: intercenter
38 study of treatment outcome in patients with complete cleft lip and palate. Part 1: introduction and
39 treatment experience. *Cleft Palate-Craniofacial Journal* 2005;42:64-8.
40
41 Semb G, Brattström V, Mølsted K, Prah-Andersen B, Zuurbier P, Rumsey N, Shaw WC. The
42 Eurocleft study: intercenter study of treatment outcome in patients with complete cleft lip and
43 palate. Part 4: relationship among outcome, patient/parent satisfaction, and the burden of care.
44 *Cleft Palate-Craniofacial Journal* 2005;42:83-92.
45
46 Shaw WC, Brattström V, Mølsted K, Prah-Andersen B, Roberts CT, Semb G. The Eurocleft study:
47 intercenter study of treatment outcome in patients with complete cleft lip and palate. Part 5:
48 discussion and conclusions. *Cleft Palate-Craniofacial Journal* 2005;42:93-8.
49
50 Tindlund RS, Rygh P. Maxillary protraction: different effects on facial morphology in unilateral and
51 bilateral cleft palate patients. *Cleft Palate-Craniofacial Journal* 1993;30:208-21.

1 Van der Heijden, P, Dijkstra PU, Stellingsma C, van der Laan BF, Korsten-Meijer AG, Goorhuis-
2 Brouwer SM. Limited evidence for the effect of presurgical nasoalveolar molding in unilateral cleft
3 on nasal symmetry: a call for unified research. *Plastic and Reconstructive Surgery* 2013;131:62e-
4 71e.

6 **Treatment Objectives and Limiting Factors**

8 American Academy of Oral and Maxillofacial Radiology. Clinical recommendations regarding use
9 of cone beam computed tomography in orthodontics. Position statement by the American
10 Academy of Oral and Maxillofacial Radiology. *Oral Surgery, Oral Medicine, Oral Pathology and*
11 *Oral Radiology* 2013;116:238-57. Erratum in: *Oral Surgery, Oral Medicine, Oral Pathology and*
12 *Oral Radiology* 2013;116:661.

14 Baird JF, Kiyak HA. The uninformed orthodontic patient and parent: treatment outcomes.
15 *American Journal of Orthodontics and Dentofacial Orthopedics* 2003;124:212–5.

17 Bayirli B, Riolo CS, Riolo ML. Treatment tactics for problems related to dentofacial discrepancies
18 in three planes of space. In: English JD, Akyalcin S, Peltomäki T, Litschel, K. *Mosby's orthodontic*
19 *review*. 2nd ed. St. Louis: Elsevier Mosby, 2015. p. 137-44.

21 Baccetti T, Franchi L, McNamara JA, Jr. Cephalometric variables predicting the long-term success
22 or failure of combined rapid maxillary expansion and facial mask therapy. *American Journal of*
23 *Orthodontics and Dentofacial Orthopedics* 2004;126:16-22.

25 Bays RA, Bouloux GF. Complications of orthognathic surgery. *Oral and Maxillofacial Surgery*
26 *Clinics of North America* 2003;15:229–42.

28 Berger JL, Pangrazio-Kulbersh V, George C, Kaczynski R. Long-term comparison of treatment
29 outcome and stability of Class II patients treated with functional appliances versus bilateral sagittal
30 split ramus osteotomy. *American Journal of Orthodontics and Dentofacial Orthopedics* 2005;
31 127:451–64.

33 Bolton WA. Disharmony in tooth size and its relation to the analysis and treatment of malocclusion.
34 *Angle Orthodontist* 1958;28:113-30.

36 Broder HL, Phillips C, Kaminetzky S. Issues in decision making: should I have orthognathic
37 surgery? *Seminars in Orthodontics* 2000;6:249–258.

39 Broadbent BH Sr, Broadbent BH Jr, Golden WH. Bolton standards of dentofacial developmental
40 growth. St. Louis: Mosby; 1975.

42 Carter L, Farman AG, Geist J, Scarfe WC, Angelopoulos C, Nair MK, Hildebolt CF, Tyndall D,
43 ShROUT M; American Academy of Oral and Maxillofacial Radiology. American Academy of Oral and
44 Maxillofacial Radiology executive opinion statement on performing and interpreting diagnostic
45 cone beam computed tomography. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology,*
46 *and Endodontics* 2008;106:561-2.

48 Chow LK, Singh B, Chiu WK, Samman N. Prevalence of postoperative complications after
49 orthognathic surgery: a 15 year review. *Journal of Oral and Maxillofacial Surgery* 2007;65:984-92.

1 Conley RS, Legan HL. Correction of severe obstructive sleep apnea with bimaxillary transverse
2 distraction osteogenesis and maxillomandibular advancement. *American Journal of Orthodontics*
3 *and Dentofacial Orthopedics* 2006;129:283–92.
4
5 Coscia G, Coscia V, Peluso V, Addabbo F. Augmented corticotomy combined with accelerated
6 orthodontic forces in Class III orthognathic patients: morphologic aspects of the mandibular
7 anterior ridge with cone-beam computer tomography. *Journal of Oral and Maxillofacial Surgery*
8 2013;71:1760.e1-9.
9
10 Gianelly AA, Arena SA, Bernstein L. A comparison of Class II treatment changes noted with the
11 light wire, edgewise, and Frankel appliances. *American Journal of Orthodontics* 1984;86:269-76.
12
13 Graber LW, Vanarsdall RL, Vig KWL. *Orthodontics: current principles and techniques*. 5th ed.
14 Philadelphia: Elsevier/Mosby; 2012.
15
16 Giblin EB, Scarpa CM. When patients refuse treatment: is it negligence if the patient elects not to
17 “follow doctor’s orders”? *AAOS Now* April 2012 Available at:
18 <http://www.aaos.org/news/aaosnow/apr12/managing6.asp>. Accessed November 5, 2015.
19
20 Greco PM. Temporomandibular disorders. In: English JD, Akyalcin S, Peltomäki T, Litschel, K.
21 *Mosby’s orthodontic review*. 2nd ed. St. Louis: Elsevier Mosby, 2015. p. 286-92.
22
23 Hoogeveen EJ, Jansma J, Ren, Y. Surgically facilitated orthodontic treatment: a systematic
24 review. *American Journal of Orthodontics and Dentofacial Orthopedics* 2014;145:S51-4.
25
26 Kulbersh R, Pangrazio-Kulbersh V. Treatment of Class II malocclusions. In: English JD, Akyalcin
27 S, Peltomäki T, Litschel, K. *Mosby’s orthodontic review*. 2nd ed. St. Louis: Elsevier Mosby, 2015. p.
28 164-85.
29
30 Little RM, Riedel RA, Artun J. An evaluation of changes in mandibular anterior alignment from 10
31 to 20 years postretention. *American Journal of Orthodontics and Dentofacial Orthopedics*
32 1988;93:423-8.
33
34 Little RM. Stability and relapse of dental arch alignment. *British Journal of Orthodontics*
35 1990;17:235-41.
36
37 Marshall SD, Southard KA, Southard TE. Phase II: nonsurgical adolescent and adult cases. In:
38 English JD, Akyalcin S, Peltomäki T, Litschel, K. *Mosby’s orthodontic review*. 2nd ed. St. Louis:
39 Elsevier Mosby, 2015. p. 206-19.
40
41 McNamara L, McNamara JA Jr. Phase I: early treatment. In: English JD, Akyalcin S, Peltomäki T,
42 Litschel, K. *Mosby’s orthodontic review*. 2nd ed. St. Louis: Elsevier Mosby, 2015. p. 145-53.
43
44 Musich DR. Assessment and description of the treatment needs of adult patients evaluated for
45 orthodontic therapy: characteristics of the solo provider group (I). *International Journal of Adult*
46 *Orthodontics and Orthognathic Surgery* 1986;1:55-67.
47
48 Musich DR. Assessment and description of the treatment needs of adult patients evaluated for
49 orthodontic therapy. II. Characteristics of the dual provider group. *International Journal of Adult*
50 *Orthodontics and Orthognathic Surgery* 1986;1:101-17.
51

1 Musich DR, Crossetti HW. Assessment and description of the treatment needs of adult patients for
2 orthodontic therapy. III. Characteristics of the multiple provider group. *International Journal of*
3 *Adult Orthodontics and Orthognathic Surgery* 1986;1:251-74.

4
5 Nance HN. The limitations of orthodontic treatment: I. Mixed dentition diagnosis and treatment.
6 *American Journal of Orthodontics and Oral Surgery* 1947;33:177-223.

7
8 Nance HN. The limitations of orthodontic treatment: II. Diagnosis and treatment in the permanent
9 dentition. *American Journal of Orthodontics and Oral Surgery* 1947;33:253-301.

10
11 Ngan P. Class III correctors. In: English JD, Akyalcin S, Peltomäki T, Litschel, K. *Mosby's*
12 *orthodontic review*. 2nd ed. St. Louis: Elsevier Mosby, 2015. p. 186-97.

13
14 Okeson JP. *Management of temporomandibular disorders and occlusion*. St. Louis:
15 Elsevier/Mosby, 2013

16
17 Pangrazio-Kulbersh V. Adult interdisciplinary orthodontic treatment. In: English JD, Akyalcin S,
18 Peltomäki T, Litschel, K. *Mosby's orthodontic review*. 2nd ed. St. Louis: Elsevier Mosby, 2015. p.
19 220-34.

20
21 Proffit WR, Fields HW, Sarver DM, Ackerman, JL. *Contemporary orthodontics*. 5th ed. St. Louis:
22 Elsevier/Mosby; 2013.

23
24 Reid KI, Green CS. Diagnosis and treatment of temporomandibular disorders: an ethical analysis
25 of current practices. *Journal of Oral Rehabilitation* 2013;40:546-61.

26
27 Roblee RD, Bolding SL, Landers JM. Surgically facilitated orthodontic therapy: a new tool for
28 optimal interdisciplinary results. *Compendium of Continuing Education in Dentistry* 2009;30:264–
29 75.

30
31 Sharpe W, Reed B, Subtelny JD, Polson A. Orthodontic relapse, apical root resorption, and crestal
32 alveolar bone levels. *American Journal of Orthodontics and Dentofacial Orthopedics* 1987;91:252-
33 8.

34
35 Strang RHW. Conditions influencing the prognosis. In: Strang RHW. *A textbook of Orthodontia*.
36 2nd ed. Philadelphia: Lea & Febiger; 1943. p. 233-5.

37
38 Tirk TM. Limitations in orthodontic treatment. *Angle Orthodontist* 1965;35:165-77.

39
40 Vaden JL, Trojan TM. Treatment planning. In: English JD, Akyalcin S, Peltomäki T, Litschel, K.
41 *Mosby's orthodontic review*. 2nd ed. St. Louis: Elsevier Mosby, 2015. p. 120-36.

42
43 van der Linden FPGM. Over de achtergronden van success en mislukking bij de behandeling van
44 angle klasse II/I-afwijkingen [Success and Failures after Treatment of Angle Class II/I Anomalies].
45 *Nederlands Tijdschrift voor Tandheelkunde* 1964;71:505-20. [Dutch]

46
47 van der Linden FPGM. *Possibilities and Limitations of Orthodontic Appliances*. Studieweek, 1965.

48 49 **Treatment Consultation and Informed Consent**

50

1 Arnett GW, Gunson MJ. Esthetic treatment planning for orthognathic surgery. *Journal of Clinical*
2 *Orthodontics* 2010;44:196-200.
3
4 Artun J. Caries and periodontal reactions associated with long-term use of different types of
5 bonded lingual retainers. *American Journal of Orthodontics* 1984;86:112-8.
6
7 Behrents RG. A treatise on the continuum of growth in the aging craniofacial skeleton: a
8 longitudinal assessment of the Bolton-Brush study participants recalled in the 1980's. Ann Arbor:
9 University of Michigan; 1984.
10
11 Copeland S, Green LJ. Root resorption in maxillary central incisors following active orthodontic
12 treatment. *American Journal of Orthodontics and Dentofacial Orthopedics* 1986;89:51-5.
13
14 Dalessandri D, Salgarello S, Dalessandri M, Lazzaroni E, Piancino M, Paganelli C, Maiorana C,
15 Santoro F. Determinants for success rates of temporary anchorage devices in orthodontics: a
16 meta-analysis (N>50). *European Journal of Orthodontics* 2014;36:303-13.
17
18 Davidson WM, Sheinis EM, Shepherd SR. Tissue reaction to orthodontic adhesives. *American*
19 *Journal of Orthodontics* 1982;82:502-7.
20
21 English JD, Akyalcin S, Peltomäki T, Litschel, K. *Mosby's orthodontic review*. 2nd ed. St. Louis:
22 Elsevier Mosby, 2015.
23
24 Geiger AM. Mucogingival problems and the movement of mandibular incisors: a clinical
25 review. *American Journal of Orthodontics* 1980;78:511-27.
26
27 Gorelick L, Geiger AM, Gwinnett AJ. Incidence of white spot formation after bonding and banding.
28 *American Journal of Orthodontics* 1982;81:93-8.
29
30 Graber LW, Vanarsdall RL, Vig KWL. *Orthodontics: current principles and techniques*. 5th ed.
31 Philadelphia: Elsevier/Mosby; 2012.
32
33 Horowitz SL, Hixon EH. Norms, classification, and treatment goals. In: Horowitz SL, Hixon EH.
34 *The nature of orthodontic diagnosis*. St. Louis; C.V. Mosby Co.; 1966. p. 325-43.
35
36 Jerrold L. Informed consent in orthodontics. *American Journal of Orthodontics and Dentofacial*
37 *Orthopedics* 1988;93:251-8.
38
39 Johnson AL. Basic principles of orthodontia. *Dental Cosmos* 1923;65:503-518.
40
41 Kersey ML, Nebbe B, Major PW. Temporomandibular joint morphology changes with mandibular
42 advancement surgery and rigid internal fixation: a systematic literature review. *Angle Orthodontist*
43 2003;73:79–85.
44
45 Kim MR, Grabber TM, Viana MA. Orthodontics and temporomandibular disorder: a meta-analysis.
46 *American Journal of Orthodontics and Dentofacial Orthopedics* 2002;121:438–46.
47
48 Langford SR, Sims MR. Upper molar root resorption because of distal movement. Report of a
49 case. *American Journal of Orthodontics* 1981;79:669-79.
50

1 Machen DE. Legal aspects of orthodontic practice: risk management concepts. *Alternative*
2 *treatment plans. American Journal of Orthodontics and Dentofacial Orthopedics* 1991;99:91-2.
3

4 Moyers RE. Standards of human occlusal development. *Craniofacial growth series No. 5* Ann
5 Arbor, MI: Center for Human Growth and Development, University of Michigan; 1976.
6

7 Musich DR Chemello PD. Orthodontic aspects of orthognathic surgery. In: Graber LW, Vanarsdall
8 RL, Vig KWL. *Orthodontics: current principles and techniques*. 5th ed. Philadelphia:
9 Elsevier/Mosby; 2012. p. 897-963.

10

11 Musich, DR. Orthodontic treatment in patients requiring jaw surgery. In: Krishnan V, Davidovitch Z.
12 *Integrated clinical orthodontics*. Chichester, West Sussex: Wiley-Blackwell; 2012. p. 332-365.
13

14 Ngan P, Wei SHY. Early treatment of Class III patients to improve facial aesthetics and predict
15 future growth. *Hong Kong Dental Journal* 2004;1:24-30. Available at:
16 [http://dentistry.hsc.wvu.edu/media/1155/early-treatment-of-class-iii-patients-to-improve-facial-](http://dentistry.hsc.wvu.edu/media/1155/early-treatment-of-class-iii-patients-to-improve-facial-aesthetics-and-predict-future-growth.pdf)
17 [aesthetics-and-predict-future-growth.pdf](http://dentistry.hsc.wvu.edu/media/1155/early-treatment-of-class-iii-patients-to-improve-facial-aesthetics-and-predict-future-growth.pdf). Accessed November 6, 2015.
18

19 Palumbo B, Cassese R, Fusetti S, Tartaro GP. Psychological aspects of orthognathic treatment.
20 *Minerva Stomatologica* 2006;55:33–42.
21

22 Popovich F, Thompson GW. Craniofacial templates for orthodontic case analysis. *American*
23 *Journal of Orthodontics* 1977;71:406-20.
24

25 Proffit WR, Fields HW, Sarver DM, Ackerman, JL. *Contemporary orthodontics*. 5th ed. St. Louis:
26 Elsevier/Mosby; 2013.
27

28 Remington DN, Joondeph DR, Artun J, Riedel, RA, Chapko MK. Long-term evaluation of root
29 resorption occurring during orthodontic treatment. *American Journal of Orthodontics and*
30 *Dentofacial Orthopedics* 1989;96:43-6.
31

32 Rinchuse DJ, Rinchuse DJ, Sosovicka MF, Robison JM, Pendleton R. Orthodontic treatment of
33 patients using bisphosphonates: a report of 2 cases. *American Journal of Orthodontics and*
34 *Dentofacial Orthopedics* 2007;131:321-6.
35

36 Rizzoli R, Burlet N, Cahall D, Delmas PD, Eriksen EF, Felsenberg D, Grbic J, Jontell M,
37 Landesberg R, Laslop A, Wollenhaupt M, Papapoulos S, Sezer O, Sprafka M, Reginster JY.
38 Osteonecrosis of the jaw and bisphosphonate treatment for osteoporosis. *Bone* 2008;42:841-7.
39

40 Roblee RD. *Interdisciplinary dentofacial therapy: a comprehensive approach to optimal patient*
41 *care*. Chicago: Quintessence; 1994.
42

43 Schiffman EL, Velly AM, Look JO, Hodges JS, Swift JQ, Decker KL, Anderson QN, Templeton RB,
44 Lenton PA, Kang W, Friction JR. Effect of four treatment strategies for temporomandibular joint
45 closed lock. *International Journal of Oral and Maxillofacial Surgery* 2014;43:217-26.
46

47 Williams AC, Shah H, Sandy JR, Travess HC. Patients' motivations for treatment and their
48 experiences of orthodontic preparation for orthognathic surgery. *Journal of Orthodontics* 2005;32:
49 191–202.
50

1 Williams DM, Bentley R, Cobourne MT, Gibilaro A, Good S, Huppa C, Matthews NS, O'Higgins E,
2 Patel S, Newton JT. Psychological characteristics of women who require orthognathic surgery:
3 comparison with untreated controls. *British Journal of Oral and Maxillofacial Surgery* 2009;47:191-
4 5.

6 Younis O, Hughes DO, Weber FN. Enamel decalcification in orthodontic treatment. *American*
7 *Journal of Orthodontics* 1979;75:678-81.

8
9 Zachrisson BU, Buyukyilmaz T. Bonding in orthodontics. In: Graber LW, Vanarsdall RL, Vig KWL.
10 *Orthodontics: current principles and techniques*. 5th ed. Philadelphia: Elsevier/Mosby; 2012. p.
11 727-84.

12 **Post Treatment Evaluation and Outcomes Assessment**

13
14
15 American Association of Orthodontists. *Guidelines for quality assessment of orthodontic care*. St.
16 Louis: American Association of Orthodontists; 1988.

17
18 Bader JD. Variation, treatment outcomes, and practice guidelines in dental practice. *Journal of*
19 *Dental Education* 1995;59:61-95.

20
21 Bailey L, Cevidanes LH, Proffit WR. Stability and predictability of orthognathic surgery. *American*
22 *Journal of Orthodontics and Dentofacial Orthopedics* 2004;236:273-7.

23
24 Bailey LJ, Dover AJ, Proffit WR. Long-term soft tissue changes after orthodontic and surgical
25 corrections of skeletal Class III malocclusions. *Angle Orthodontist* 2007;77:389–96.

26
27 Bailey LJ, Phillips C, Proffit WR. Long-term outcome of surgical Class III correction as a function of
28 age at surgery. *American Journal of Orthodontics and Dentofacial Orthopedics* 2008;133:365–70.

29
30 Bishara SE. Impacted maxillary canines: a review. *American Journal of Orthodontics and*
31 *Dentofacial Orthopedics* 1992;101:159-71.

32
33 Boyd RL. Two-year longitudinal study of a peroxide-fluoride rinse on decalcification in adolescent
34 orthodontic patients. *Journal of Clinical Dentistry* 1992;3:83-7.

35
36 Chateau M, Demoge PH. Evaluation of long term results of orthodontic therapy. *International*
37 *Dental Journal* 1961;11:29-46.

38
39 Chen SS, Greenlee GM, Kim JE, Smith CL, Huang GJ. Systematic review of self-ligating brackets.
40 *American Journal of Orthodontics and Dentofacial Orthopedics* 2010;137:726.e1-18.

41
42 Chen Y, Kyung HM, Zhao WT, Yu WJ. Critical factors for the success of orthodontic mini-implants:
43 a systematic review. *American Journal of Orthodontics and Dentofacial Orthopedics*
44 2009;135:284-91.

45
46 Scott Conley R, Jernigan C. Soft tissue changes after upper premolar extraction in Class II
47 camouflage therapy. *Angle Orthodontist* 2006;76:59–65.

48
49 Dyken RA, Sadowsky PL, Hurst D. Orthodontic outcomes assessment using the peer assessment
50 rating index. *Angle Orthodontist* 2001;71:164-9.

51

1 Goto S, Boyd RL, Nielsen L, Iizuka T. Long-term followup of orthodontic treatment of a patient with
2 maxillary protrusion, severe deep overbite and thumb-sucking. *Angle Orthodontist* 1994;64:7-12.
3

4 Graber TM. Postmortems in posttreatment adjustment. *American Journal of Orthodontics*
5 1966;52:331-52.
6

7 Huang GJ, Roloff-Chiang B, Mills BE, Shalchi S, Spiekerman C, Korpak AM, Starrett JL, Greenlee
8 GM, Drangsholt RJ, Matunas JC. Effectiveness of MI Paste Plus and PreviDent fluoride varnish for
9 treatment of white spot lesions: a randomized controlled trial. *American Journal of Orthodontics*
10 and *Dentofacial Orthopedics* 2013;143:31-41.
11

12 Kim MR, Graber TM, Viana MA. Orthodontics and temporomandibular disorder: a meta-analysis.
13 *American Journal of Orthodontics and Dentofacial Orthopedics* 2002;121:438-46.
14

15 Korkhaus G (moderator). Posttreatment appraisal of orthodontic results. *European Orthodontic*
16 *Society Transactions* 1961;73-97.
17

18 Kim SJ, Kim MR, Shin SW, Chun YS, Kim EJ. Evaluation on the psychosocial status of
19 orthognathic surgery patients. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and*
20 *Endodontology* 2009;108):828-32.
21

22 Little RM, Riedel RA, Engst ED. Serial extraction of first premolars-postretention evaluation of
23 stability and relapse. *Angle Orthodontist* 1990;60:255-62.
24

25 Little RM, Riedel RA, Stein A. Mandibular arch length increase during the mixed dentition:
26 postretention evaluation of stability and relapse. *American Journal of Orthodontics and Dentofacial*
27 *Orthopedics* 1990;97:393-404.
28

29 Little RM, Riedel RA. Postretention evaluation of stability and relapse-mandibular arches with
30 generalized spacing. *American Journal of Orthodontics and Dentofacial Orthopedics* 1989;95:37-
31 41.
32

33 McReynolds DC, Little RM. Mandibular second premolar extraction-postretention evaluation of
34 stability and relapse. *Angle Orthodontist* 1991;61:133-44.
35

36 Marshall SD, Currier GF, Hatch NE, Huang GJ, Nah HD, Owens SE, Shroff B, Southard TE, Suri
37 L, Turpin DL. Ask us. Self-ligating bracket claims. *American Journal of Orthodontics and*
38 *Dentofacial Orthopedics* 2010;138:128-31.
39

40 O'Brien K, Wright J, Conboy F, Appelbe P, Davies L, Connolly I, Mitchell L, Littlewood S, Mandall
41 N, Lewis D, Sandler J, Hammond M, Chadwick S, O'Neill J, McDade C, Oskouei M,
42 Thiruvengkatachari B, Read M, Robinson S, Birnie D, Murray A, Shaw I, Harradine N, Worthington
43 H. Early treatment for Class II Division 1 malocclusion with the Twin-block appliance: a multi-
44 center, randomized, controlled trial. *American Journal of Orthodontics and Dentofacial*
45 *Orthopedics* 2009;135:573-79.
46

47 Ormiston JP, Huang GJ, Little RM, Decker JD, Seuk GD. Retrospective analysis of long-term
48 stable and unstable treatment outcomes. *American Journal of Orthodontics and Dentofacial*
49 *Orthopedics* 2005;128:568-74.
50

- 1 Pennsylvania Dental Association. Quality assessment guidelines. Harrisburg, PA: Pennsylvania
2 Dental Association; 1993.
- 3
- 4 Riedel RA, Little RM, Bui TD. Mandibular incisor extraction-postretention evaluation of stability and
5 relapse. *Angle Orthodontist* 1992;62:103-16.
- 6
- 7 Sadowsky C, Theisen TA, Sakols EI. Orthodontic treatment and temporomandibular joint sounds-a
8 longitudinal study. *American Journal of Orthodontics and Dentofacial Orthopedics* 1991;99:441-7.
- 9
- 10 Segal GR, Schiffman PH, Tuncay OC. Meta analysis of the treatment-related factors of external
11 apical root resorption. *Orthodontics and Craniofacial Research* 2004;7:71-8.
- 12
- 13 Sforza C, Laino A, D'allessio R, Grandi G, Tartaglia GM, Ferrario VF. Soft-tissue facial
14 characteristics of attractive and normal adolescent boys and girls. *Angle Orthodontist*
15 2008;78:799–807.
- 16
- 17 Stellzig-Eisenhauer A, Lux CJ, Schuster G. Treatment decision in adult patients with Class III
18 malocclusion: orthodontic therapy or orthognathic surgery? *American Journal of Orthodontics and*
19 *Dentofacial Orthopedics* 2002;122:27–38.
- 20
- 21 Suri L, Taneja P. Surgically assisted rapid palatal expansion: a literature review. *American Journal*
22 *of Orthodontics and Dentofacial Orthopedics* 2008;133:290-302.
- 23
- 24 Vastardis H. The genetics of human tooth agenesis: new discoveries for understanding dental
25 anomalies. *American Journal of Orthodontics and Dentofacial Orthopedics* 2000;177:650-56.
- 26
- 27 Wade DB. Outcomes assessed by orthodontic programs. *American Journal of Orthodontics and*
28 *Dentofacial Orthopedics* 1994;106:109.
- 29
- 30 Weltman B, Vig KW, Fields HW, Shanker S, Kaizar EE. Root resorption associated with
31 orthodontic tooth movement: a systematic review. *American Journal of Orthodontics and*
32 *Dentofacial Orthopedics* 2010;137:462-76.
- 33
- 34 Wolford LM, Chemello PD, Hilliard F. Occlusal plane alteration in orthognathic surgery - Part I:
35 Effects of function and esthetics. *American Journal of Orthodontics and Dentofacial Orthopedics*
36 1994;106:304–16.
- 37
- 38 **Retention**
- 39
- 40 Akyalcin S, Kapadia H, English JD. Retention and relapse in orthodontics. In: English JD, Akyalcin
41 S, Peltomäki T, Litschel, K. *Mosby's orthodontic review*. 2nd ed. St. Louis: Elsevier Mosby, 2015. p.
42 293-301.
- 43
- 44 Behrents RG. A treatise on the continuum of growth in the aging craniofacial skeleton. [thesis] Ann
45 Arbor, MI: University of Michigan; 1984.
- 46
- 47 Bjerling R, Birkeland K, Vandevska-Radunovic V. Anterior tooth alignment: A comparison of
48 orthodontic retention regimens 5 years posttreatment. *Angle Orthodontist* 2015;85:353-9.
- 49
- 50 Bondevik O. A longitudinal study of dental arches and occlusal changes in adults from 21-33, and
51 33 to 43 years of age. *Journal of Orofacial Orthopedics* 2015;76:79-89.

1 Goldberg AI, Behrents RG, Oliver DR, Buschang PH. Facial divergence and mandibular crowding
2 in treated subjects. *Angle Orthodontist* 2013;83:381-8.
3

4 Kaplan H. The logic of modern retention procedures. *American Journal of Orthodontics and*
5 *Dentofacial Orthopedics* 1988;93:325-40.
6

7 Kurt G, Altug-Ataç AT, Ataç MS, Karasu HA. Stability of surgically assisted rapid maxillary
8 expansion and orthopedic maxillary expansion after 3 years' follow-up. *Angle Orthodontist* 2010;
9 80:425-31.
10

11 Lenz GJ, Woods MG. Incisal changes and orthodontic stability. *Angle Orthodontist* 1999;69:424-
12 32.
13

14 Little RM, Riedel RA, Artun J. An evaluation of changes in mandibular anterior alignment from 10
15 to 20 years postretention. *American Journal of Orthodontics and Dentofacial Orthopedics*
16 1988;93:423-8.
17

18 Little RM, Wallen TR, Riedel RA. Stability and relapse of mandibular anterior alignment-first
19 premolar extraction cases treated by traditional edgewise orthodontics. *American Journal of*
20 *Orthodontics* 1981;80:349-65.
21

22 Lopez-Gavito G, Wallen TR, Little RM, Joondeph DR. Anterior open-bite malocclusion: a
23 longitudinal 10-year postretention evaluation of orthodontically treated patients. *American Journal*
24 *of Orthodontics* 1985;87:175-86.
25

26 Maia NG, Normando D, Maia FA, Ferreira MA, do Socorro Costa Feitosa Alves M. Factors
27 associated with long-term patient satisfaction. *Angle Orthodontist* 2010;80:1155-8.
28

29 Makki L, Ferguson DJ, Wilcko MT, Wilcko WM, Bjerklin K, Stapelberg R, Al-Mulla A. Mandibular
30 irregularity index stability following alveolar corticotomy and grafting: a 10 year preliminary study.
31 *Angle Orthodontist* 2015;85:743-9.
32

33 Mucedero M, Coviello A, Baccetti T, Franchi L, Cozza P. Stability factors after double-jaw surgery
34 in Class III malocclusion. A systematic review. *Angle Orthodontist* 2008;78:1141-52.
35

36 Palma JC, Tejedor-Sanz N, Oteo MD, Alarcón JA. Long-term stability of rapid maxillary expansion
37 combined with chincup protraction followed by fixed appliances. *Angle Orthodontist* 2015;85:270-
38 7.
39

40 Proffit WR, Bailey LJ, Phillips C, Turvey TA. Long-term stability of surgical open-bite correction by
41 Le Fort I osteotomy. *Angle Orthodontist* 2000;70:112-7.
42

43 Reitan K. Tissue rearrangement during retention of orthodontically rotated teeth. *Angle*
44 *Orthodontist* 1959;29:105-13.
45

46 Reitan K. Principles of retention and avoidance of posttreatment relapse. *American Journal of*
47 *Orthodontics* 1969;55:776-90.
48

49 Solomon MJ, English JD, Magness WB, McKee CJ. Long-term stability of lip bumper therapy
50 followed by fixed appliances. *Angle Orthodontist* 2006;76:36-42.
51

1 Sondhi A, Cleall JF, BeGole EA. Dimensional changes in the dental arches of orthodontically
2 treated cases. American Journal of Orthodontics 1980;77:60-74.

3
4 Wiedel AP, Bondemark L. Stability of anterior crossbite correction: a randomized controlled trial
5 with a 2-year follow-up. Angle Orthodontist 2015;85:189-95.

6
7 Zachrisson BU. Adult retention: a new approach. In: Graber LW, Graber TM. Orthodontics, state of
8 the art, essence of the science. St. Louis: Mosby; 1986. p. 310-27.

9 10 **Record Keeping**

11
12 American Association of Orthodontists. Orthodontics a patient education guide. St. Louis:
13 American Association of Orthodontists; 1991.

14
15 Eash C. Personnel file and recordkeeping. American Journal of Orthodontics and Dentofacial
16 Orthopedics 1994;105:610-1.

17
18 Jerrold L. Dental records and record keeping. American Journal of Orthodontics and Dentofacial
19 Orthopedics 1993;104:98-9.

20
21 Jerrold L. Litigation and Legislation. Destroying records. American Journal of Orthodontics and
22 Dentofacial Orthopedics 2015;148:184-6.

23
24 Jerrold L. Litigation and Legislation. What record is playing? American Journal of Orthodontics and
25 Dentofacial Orthopedics 2015;147:627-31.

26
27 Jerrold L. Litigation, legislation, and ethics. Dental records as evidence. American Journal of
28 Orthodontics and Dentofacial Orthopedics 2000;118:241-2.

29
30 Jerrold L. Litigation, legislation, and ethics. Posttreatment records. American Journal of
31 Orthodontics and Dentofacial Orthopedics 2008;133:124-6.

32
33 Machen DE. Legal aspects of orthodontic practice: risk management concepts. Excellent
34 diagnostic informed consent practice and record keeping make a difference. American Journal of
35 Orthodontics and Dentofacial Orthopedics 1990;98:381-2.

36
37 Morin DR. The patient's records and the defense of dental malpractice claims. American Journal
38 of Orthodontics and Dentofacial Orthopedics 1992;102:569-70.

39
40 Speidel TM, Jerrold L. Litigation, legislation, and ethics. Record keeping to avoid or defend
41 lawsuits: a defense attorney's perspective. American Journal of Orthodontics and Dentofacial
42 Orthopedics 2004;125:754-6.

43 44 **Transfer of Orthodontic Patients**

45
46 American Association of Orthodontists. American Association of Orthodontists bylaws and
47 principles of ethics. St. Louis: American Association of Orthodontists; 1994.

48
49 American Association of Orthodontists. Guidelines for transfer of orthodontic cases. St. Louis:
50 American Association of Orthodontists; 1993.

- 1 Jerrold L. Litigation and legislation: problems of our own making: a view from both sides of the
- 2 coin. American Journal of Orthodontics and Dentofacial Orthopedics 2012;142:734-8.
- 3
- 4 Jerrold L. Litigation and Legislation. Transfer tragedies. American Journal of Orthodontics and
- 5 Dentofacial Orthopedics 2015;147:788-90.
- 6